

BALANCED SIMPLICES

JONATHAN CHAPPELON

ABSTRACT. An additive cellular automaton is a linear map on the set of infinite multi-dimensional arrays of elements in a finite cyclic group $\mathbb{Z}/m\mathbb{Z}$. In this paper, we consider simplices appearing in the orbits generated from arithmetic arrays by additive cellular automata. We prove that they constitute a source of balanced simplices, that are simplices containing all the elements of $\mathbb{Z}/m\mathbb{Z}$ with the same multiplicity. For any additive cellular automaton of dimension 1 or higher, the existence of infinitely many balanced simplices of $\mathbb{Z}/m\mathbb{Z}$ appearing in such orbits is shown, and this, for an infinite number of values m . The special case of the Pascal cellular automata, the cellular automata generating the multidimensional simplices of Pascal, is studied in detail.

1. INTRODUCTION

Let n and m be positive integers. Throughout this paper, n will denote the dimension of the objects studied and m the order of the finite cyclic group $\mathbb{Z}/m\mathbb{Z}$. For any integers a and b such that $a < b$, we denote by $[a, b]$ the set of the integers between a and b , that is, $[a, b] := \{a, a + 1, \dots, b\}$ and by $[a, b]^n$ the Cartesian product of n copies of $[a, b]$. For any n -tuple of elements u , we denote by u_i its i th component for all $i \in [1, n]$, that is, $u = (u_1, \dots, u_n)$. For two n -tuples u and v and an integer λ , we consider the sum $u + v := (u_1 + v_1, \dots, u_n + v_n)$, the product $u \cdot v := (u_1 v_1, \dots, u_n v_n)$ and the scalar product $\lambda u := (\lambda u_1, \dots, \lambda u_n)$.

Definition 1.1 (ACA). Let r be a non-negative integer and let $W = (w_j)_{j \in [-r, r]^n}$ be an n -dimensional array of integers of size $(2r + 1)^n$. The *additive cellular automaton* (ACA for short) over $\mathbb{Z}/m\mathbb{Z}$ associated with W is the map ∂ which assigns, to every n -dimensional infinite array of $\mathbb{Z}/m\mathbb{Z}$, a new array by a linear transformation whose coefficients are those of W . More precisely, the map ∂ is defined by

$$\partial((a_i)_{i \in \mathbb{Z}^n}) = \left(\sum_{j \in [-r, r]^n} w_j a_{i+j} \right)_{i \in \mathbb{Z}^n},$$

for all arrays $(a_i)_{i \in \mathbb{Z}^n}$ of elements in $\mathbb{Z}/m\mathbb{Z}$. We say that ∂ is of *dimension* $n \geq 1$ and of *weight* W with *radius* $r \geq 0$.

Definition 1.2 (Orbit). Let $A = (a_i)_{i \in \mathbb{Z}^n}$ be an infinite array of $\mathbb{Z}/m\mathbb{Z}$ of dimension n . The *orbit* $\mathcal{O}(A)$ generated from A by the ACA ∂ is the collection of all the n -dimensional arrays obtained from A by successive applications of ∂ , that is,

$$\mathcal{O}(A) := \{\partial^j(A) \mid j \in \mathbb{N}\},$$

where ∂^j is recursively defined by $\partial^j(A) = \partial(\partial^{j-1}(A))$ for all $j \geq 1$ and $\partial^0(A) = A$. The orbit $\mathcal{O}(A)$ can also be seen as the $(n + 1)$ -dimensional array $(a_{i,j})_{(i,j) \in \mathbb{Z}^n \times \mathbb{N}}$ of $\mathbb{Z}/m\mathbb{Z}$ whose j th row $R_j := (a_{i,j})_{i \in \mathbb{Z}^n}$ corresponds to $\partial^j(A)$, for all $j \in \mathbb{N}$.

Date: February 19, 2014.

2010 Mathematics Subject Classification. 05B30, 11B75, 05A05, 11B50, 11A99.

Key words and phrases. Balanced simplices, additive cellular automaton, Pascal cellular automaton, Steinhaus triangles, Pascal triangles, Molluzzo's problem.

$a_{0,0}$	$a_{1,0}$	2	3	4	0	1	2	3	4	0	1	2	3	4	0
$a_{0,1}$	$a_{1,1}$	2	1	0	4	3	2	1	0	4	3	2	1	0	4
2	3	4	0	1	2	3	4	0	1	2	3	4	0	1	2
2	1	0	4	3	2	1	0	4	3	2	1	0	4	3	2
4	0	1	2	3	4	0	1	2	3	4	0	1	2	3	4
0	4	3	2	1	0	4	3	2	1	0	4	3	2	1	0
1	2	3	4	0	1	2	3	4	0	1	2	3	4	0	1
4	3	2	1	0	4	3	2	1	0	4	3	2	1	0	4
2	3	4	0	1	2	3	4	0	1	2	3	4	0	1	2
2	1	0	4	3	2	1	0	4	3	2	1	0	4	3	2
4	0	1	2	3	4	0	1	2	3	4	0	1	2	3	4
0	4	3	2	1	0	4	3	2	1	0	4	3	2	1	0
1	2	3	4	0	1	2	3	4	0	1	2	3	4	0	1
4	3	2	1	0	4	3	2	1	0	4	3	2	1	0	4
2	3	4	0	1	2	3	4	0	1	2	3	4	0	1	2
2	1	0	4	3	2	1	0	4	3	2	1	0	4	3	2

FIGURE 1. Example of triangles $\Delta((2, 2), ++, 5)$, $\Delta((2, 13), +-, 5)$, $\Delta((13, 2), -+, 5)$ and $\Delta((13, 13), --, 5)$ appearing in an orbit $\mathcal{O}(A) = (a_{i,j})_{(i,j) \in \mathbb{Z} \times \mathbb{N}}$ of $\mathbb{Z}/5\mathbb{Z}$ generated by the ACA of weight $W = (2, 1, 1)$.




FIGURE 2. The eight possible orientations of a tetrahedron

Definition 1.3 (Simplices). Let $A = (a_i)_{i \in \mathbb{Z}^n}$ be an infinite array of $\mathbb{Z}/m\mathbb{Z}$ of dimension n . Let $\varepsilon \in \{-1, 1\}^n$ and let s be a positive integer. The *simplex* of size s , with orientation ε and whose principal vertex is at the coordinates $j \in \mathbb{Z}^n$ in A , is the multiset of $\mathbb{Z}/m\mathbb{Z}$ defined and denoted by

$$\Delta(j, \varepsilon, s) := \{a_{j+\varepsilon \cdot k} \mid k \in \mathbb{N}^n \text{ such that } k_1 + \cdots + k_n \leq s - 1\}.$$

For $n = 2$ and $n = 3$, it is called a triangle and a tetrahedron, respectively.

In this paper, we mainly consider simplices of dimension n appearing in the orbit generated from an infinite array of $\mathbb{Z}/m\mathbb{Z}$ by an ACA of dimension $n - 1$. Examples of triangles, for the four possible orientations in dimension $n = 2$, are depicted in Figure 1.

In Figure 2, for dimension $n = 3$, the eight possible orientations of a tetrahedra are represented.

Let M be a *multiset* of $\mathbb{Z}/m\mathbb{Z}$, that is a set where each element of $\mathbb{Z}/m\mathbb{Z}$ can appear more than once. The *multiplicity function* associated to M is the integer-valued function $\mathbf{m}_M : \mathbb{Z}/m\mathbb{Z} \mapsto \mathbb{N}$, which assigns to each element of $\mathbb{Z}/m\mathbb{Z}$ its multiplicity in M . The *cardinality* of M , denoted by $|M|$, is the number of elements constituting M , counted with multiplicity, that is, $|M| = \sum_{x \in \mathbb{Z}/m\mathbb{Z}} \mathbf{m}_M(x)$.

Definition 1.4 (Balanced multisets of $\mathbb{Z}/m\mathbb{Z}$). A multiset of $\mathbb{Z}/m\mathbb{Z}$ is said to be *balanced* if it contains all the elements of $\mathbb{Z}/m\mathbb{Z}$ with the same multiplicity, i.e., if its associated multiplicity function \mathbf{m}_M is constant on $\mathbb{Z}/m\mathbb{Z}$, equal to $\frac{1}{m}|M|$.

The goal of this paper is to prove the existence of balanced simplices appearing in certain orbits generated by ACA. Sufficient conditions for obtaining this result will be detailed throughout this paper. This notion of balanced simplices generated by ACA essentially appears in the literature in the case of the Pascal cellular automaton of dimension 1.

Definition 1.5 (PCA_n). The *Pascal cellular automaton* of dimension n is the ACA of radius $r = 1$ and whose weight array $W = (w_i)_{i \in [-1,1]^n}$ is defined by

$$w_i = \begin{cases} 1 & \text{if } i \in \{0_{\mathbb{Z}^n}, -e_1, -e_2, \dots, -e_n\}, \\ 0 & \text{otherwise,} \end{cases}$$

where (e_1, e_2, \dots, e_n) is the canonical basis of the vector space \mathbb{Z}^n . It is denoted by PCA_n .

For instance, $W = (1, 1, 0)$ for PCA_1 and $W = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ for PCA_2 .

Remark 1.6. Let $A = (a_i)_{i \in \mathbb{Z}^{n-1}}$ be the $(n-1)$ -dimensional array of $\mathbb{Z}/m\mathbb{Z}$ defined by $a_i = 1$ for $i = 0_{\mathbb{Z}^n}$ and $a_i = 0$ otherwise. If $(a_{i,j})_{(i,j) \in \mathbb{Z}^{n-1} \times \mathbb{N}}$ is the orbit $\mathcal{O}(A)$ generated by the PCA_{n-1} , then $a_{i,j}$ is the multinomial coefficient

$$a_{i,j} = \binom{j}{i_1, \dots, i_{n-1}, j - \sum_{k=1}^{n-1} i_k} = \frac{j!}{i_1! \cdots i_{n-1}! (j - \sum_{k=1}^{n-1} i_k)!}$$

for all $i \in \mathbb{N}^{n-1}$ such that $i_1 + \cdots + i_{n-1} \leq j$, and $a_{i,j} = 0$ otherwise. Thus, we retrieve the coefficients of the Pascal n -simplex modulo m . This is the reason why this specific ACA is called the Pascal cellular automaton.

Let $A = (a_i)_{i \in \mathbb{Z}}$ be a doubly infinite sequence of $\mathbb{Z}/m\mathbb{Z}$. We consider the orbit generated from A by PCA_1 , i.e., the infinite array $\mathcal{O}(A) = (a_{i,j})_{(i,j) \in \mathbb{Z} \times \mathbb{N}}$ defined by $a_{i,0} = a_i$ for all $i \in \mathbb{Z}$ and

$$a_{i,j} = a_{i-1,j-1} + a_{i,j-1}$$

for all $(i,j) \in \mathbb{Z} \times \mathbb{N}^*$. The $(-+)$ -triangles and $(+-)$ -triangles appearing in a such orbit are known as *Steinhaus triangles* and *generalized Pascal triangles* modulo m , respectively. The name of Steinhaus triangle is due to Hugo Steinhaus himself in [26], where he proposed this construction in the binary case $m = 2$. He posed the following problem, as an unsolved problem.

Problem 1.7 (Steinhaus [26]). For every positive integer s such that $s \equiv 0$ or $3 \pmod{4}$, does there exist a Steinhaus triangle of size s in $\mathbb{Z}/2\mathbb{Z}$ containing as many 0's as 1's?

Remark that the condition on the size s of a balanced Steinhaus triangle in $\mathbb{Z}/2\mathbb{Z}$ is obviously a necessary condition because the number of elements of a such triangle, that is $\binom{s+1}{2}$, is even if and only if $s \equiv 0$ or $3 \pmod{4}$. A positive solution to this problem

appeared in the literature for the first time in [23], where the author gave, for every $s \equiv 0$ or $3 \pmod{4}$, an explicit construction of a balanced Steinhaus triangle of size s in $\mathbb{Z}/2\mathbb{Z}$. More recently, several other constructions of balanced binary Steinhaus triangles have been obtained by considering sequences with additional properties such as strongly balanced [20], symmetric and antisymmetric [21], or zero-sum sequences [22].

The number of elements $1 \in \mathbb{Z}/2\mathbb{Z}$ appearing in binary Steinhaus triangles and in binary generalized Pascal triangles are studied in [6] and in [24], respectively. Such binary triangles having certain geometric properties are studied in [3, 5]. A binary Steinhaus triangle can also be considered as the upper triangular part of the adjacency matrix of a finite graph. These undirected graphs are called Steinhaus graphs in [25]. A classical problem on Steinhaus graphs is to study those having certain graphical properties such as bipartition [7, 15, 18], planarity [17] or regularity [1, 2, 10, 14]. A survey on Steinhaus graph can be found in [16].

Problem 1.7 on balanced binary Steinhaus triangles was generalized for any positive integers m by Molluzzo in [25].

Problem 1.8 (Molluzzo [25]). Let m be a positive integer. For every positive integer s such that $\binom{s+1}{2}$ is divisible by m , does there exist a Steinhaus triangle of size s containing all the elements of $\mathbb{Z}/m\mathbb{Z}$ with the same multiplicity?

Since, this problem has been positively solved, by constructive approaches, for small values of m : for $m \in \{3, 5\}$ in [4], for $m \in \{3, 5, 7\}$ in [8], for $m = 4$ in [13]. First counter-examples appeared in [8], where the author proved that there does not exist balanced Steinhaus triangles of size 5 in $\mathbb{Z}/15\mathbb{Z}$ and of size 6 in $\mathbb{Z}/21\mathbb{Z}$. Nevertheless, this problem can be positively answered for an infinite number of values m . Indeed, as showed in [8, 9], there exist balanced Steinhaus triangles, for all the possible sizes, in the case where m is a power of 3. More precisely, the author obtained the following result.

Theorem 1.9 (Chappelton [8, 9]). *Let m be an odd number and let a and d be in $\mathbb{Z}/m\mathbb{Z}$ such that d is invertible. The Steinhaus triangle, of size s , whose first row is the arithmetic progression $(a, a + d, a + 2d, \dots, a + (s - 1)d)$ is balanced in $\mathbb{Z}/m\mathbb{Z}$, for all $s \equiv 0$ or $-1 \pmod{\text{ord}_m(2^m)m}$, where $\text{ord}_m(2^m)$ is the multiplicative order of 2^m modulo m , i.e., the order of 2^m in the group of invertibles $(\mathbb{Z}/m\mathbb{Z})^*$.*

In particular, since $\text{ord}_m(2^m) = 2$ for all $m = 3^k$, where k is a positive integer, it follows from Theorem 1.9 that there exist balanced Steinhaus triangles of size s in $\mathbb{Z}/3^k\mathbb{Z}$ for all $s \equiv 0$ or $-1 \pmod{2 \cdot 3^k}$. This result can be refined by considering Steinhaus triangles whose first row has the additional property to be antisymmetric. Thus, the author obtained a positive answer to the Molluzzo problem for all $m = 3^k$. Even if the Molluzzo problem is not completely solved for the other odd values of m , we know from Theorem 1.9 that there exist infinitely many balanced Steinhaus triangles in every $\mathbb{Z}/m\mathbb{Z}$ with m odd. This weak version of the Molluzzo problem was posed in [13].

Problem 1.10 (Weak Molluzzo problem). Let m be a positive integer. Does there exist infinitely many balanced Steinhaus triangles of $\mathbb{Z}/m\mathbb{Z}$?

Problem 1.10 is thus solved for the odd numbers m . For the even values, the cases $m = 2$ and $m = 4$ come from the solutions of Problem 1.8 and a solution will appear in [19] for $m \in \{6, 8, 10\}$. This problem is completely open for the even numbers $m \geq 12$. Similar problems of determining the existence of balanced structures can be considered for other shapes. In [12], the author not only considers balanced Steinhaus triangles, but also balanced generalized Pascal triangles, trapezoids or lozenges. In particular, for Steinhaus triangles and generalized Pascal triangles, the following result is proved.

0	1	2	3	4	3	4	0	1	1	2	3	4	0	2		
2	3	4	0	0	1	2	3	4	0			1				
4	0	1			2	3			0							
1	2					4										
3																

FIGURE 3. The arithmetic tetrahedron $AS(0, (1, 2, 3), 5)$ in $\mathbb{Z}/5\mathbb{Z}$

Theorem 1.11 (Chappelon, [12]). *Let m be an odd number. For all $s \equiv 0 \pmod{m}$ and $s \equiv -1 \pmod{3m}$, there exist Steinhaus triangles and generalized Pascal triangles of size s , which are balanced in $\mathbb{Z}/m\mathbb{Z}$.*

In fact, the author proves that the orbit generated by PCA_1 from the following sequence

$$(\dots, 0, -1, 1, 1, -3, 2, 2, -5, 3, 3, -7, 4, 4, -9, 5, 5, \dots),$$

which is an interlacing of three arithmetic progressions, contains all the balanced triangles whose existence is announced in Theorem 1.11 and this, for every odd number m . The main ingredient in the proof of this result is an elementary object that the author called a doubly arithmetic triangle in [12] and that we simply call an arithmetic triangle here in this paper. An arithmetic triangle with common differences d_1 and d_2 is a triangle of elements in $\mathbb{Z}/m\mathbb{Z}$ whose rows and columns are arithmetic progressions with the same common differences, d_1 for the rows and d_2 for the columns. The interest of this structure is that this is very often balanced. In this paper, we consider a generalization in higher dimensions of arithmetic triangles.

Definition 1.12 (Arithmetic arrays and simplices). Let n and m be positive integers. Let $A = (a_i)_{i \in \mathbb{Z}^n}$ be an array of $\mathbb{Z}/m\mathbb{Z}$. The array A is said to be *arithmetic* with first element a and with common difference $d = (d_1, \dots, d_n) \in (\mathbb{Z}/m\mathbb{Z})^n$ if

$$a_i = a + i_1 d_1 + \dots + i_n d_n,$$

for all $i = (i_1, \dots, i_n) \in \mathbb{Z}^n$. The arithmetic array with first element $a \in \mathbb{Z}/m\mathbb{Z}$ and with common difference $d \in (\mathbb{Z}/m\mathbb{Z})^n$ is denoted by $AA(a, d)$. The *arithmetic simplex* of size s , with first element $a \in \mathbb{Z}/m\mathbb{Z}$ and with common difference $d = (d_1, \dots, d_n) \in (\mathbb{Z}/m\mathbb{Z})^n$, is the simplex $\triangle(0_{\mathbb{Z}^n}, + \dots +, s)$ appearing in the array $AA(a, d) = (a_i)_{i \in \mathbb{Z}^n}$ and is denoted by $AS(a, d, s)$, that is,

$$AS(a, d, s) = \{a + i_1 d_1 + \dots + i_n d_n \mid i \in \mathbb{N}^n \text{ such that } i_1 + \dots + i_n \leq s - 1\}.$$

For $n = 1$, the arithmetic progression $AS(a, d, s)$ is also denoted by $AP(a, d, s)$.

Remark 1.13. The following multiset identities hold for arithmetic simplices :

$$\begin{aligned} AS(a, (d_1, \dots, d_n), s) &= AS(a + (s-1)d_1, (-d_1, d_2 - d_1, \dots, d_n - d_1), s) \\ &= AS(a + (s-1)d_2, (d_1 - d_2, -d_2, d_3 - d_2, \dots, d_n - d_2), s) \\ &\dots\dots\dots \\ &= AS(a + (s-1)d_n, (d_1 - d_n, \dots, d_{n-1} - d_n, -d_n), s), \end{aligned}$$

and $AS(a, (d_1, \dots, d_n), s) = AS(a, (d_{\pi(1)}, \dots, d_{\pi(n)}), s)$, where π is a permutation of $[1, n]$.

For example, the arithmetic tetrahedron $AS(0, (1, 2, 3), 5)$ of $\mathbb{Z}/5\mathbb{Z}$ is depicted in Figure 3. The successive rows of this tetrahedron are the arithmetic triangles $AS(0, (1, 2), 5)$, $AS(3, (1, 2), 4)$, $AS(1, (1, 2), 3)$, $AS(4, (1, 2), 2)$ and $AS(2, (1, 2), 1)$ of $\mathbb{Z}/5\mathbb{Z}$.

Return now to the general case of an ACA of dimension $n - 1$, with a weight array $W = (w_j)_{j \in [-r, r]^{n-1}}$ of radius $r \in \mathbb{N}$. Let us denote

$$\sigma := \sum_{j \in [-r, r]^{n-1}} w_j \quad \text{and} \quad \sigma_k := \sum_{j \in [-r, r]^{n-1}} j_k w_j, \quad \text{for all } k \in [1, n-1].$$

For any integers a and b , we denote by $\gcd(a, b)$ and $\text{lcm}(a, b)$ the greatest common divisor and the least common multiplier of a and b , respectively. Let $x \in \mathbb{Z}/m\mathbb{Z}$. We also denote by $\gcd(x, m)$ the greatest common divisor of m and any representant of the residue class x .

Using properties of arithmetic simplices, the following theorem, which is the main result of this paper, will be proved.

Theorem 1.14. *Let $n \geq 2$ and m be two positive integers such that $\gcd(m, n!) = 1$. Suppose that σ is invertible modulo m . Let $a \in \mathbb{Z}/m\mathbb{Z}$, $d = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$ and $\varepsilon \in \{-1, 1\}^n$ such that d_i , for all $1 \leq i \leq n$, and $\varepsilon_j d_j - \varepsilon_i d_i$, for all $1 \leq i < j \leq n$, are invertible, where $d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k$. Then, in the orbit $\mathcal{O}(\text{AA}(a, d))$, every n -simplex with orientation ε and of size s is balanced, for all $s \equiv -t \pmod{\text{lcm}(\text{ord}_m(\sigma), m)}$, where $t \in [0, n-1]$.*

Remark 1.15. Obviously, $\text{lcm}(\text{ord}_m(\sigma), m) = \text{ord}_m(\sigma^m)m$. A complete study of this arithmetic function can be found in [11].

For $n = 2$, m odd and $W = (1, 1, 0)$, the weight sequence of PCA_1 , we retrieve Theorem 1.9. Indeed, in this case, we have $\sigma = 2$, $\sigma_1 = -1$, $d_1 = d$, $d_2 = \sigma^{-1} \sigma_1 d_1 = -2^{-1}d$ and $\varepsilon_1 d_1 - \varepsilon_2 d_2 = \pm 2^{-1}d$ for $\varepsilon = (\pm 1, \mp 1)$, which are invertibles of $\mathbb{Z}/m\mathbb{Z}$.

In the special case of PCA_{n-1} , Theorem 1.14 gives a positive answer to the equivalent problem of the weak Molluzzo problem, in higher dimensions, for an infinite number of values m .

Corollary 1.16. *Let $n \geq 2$ be a positive integers. For every positive integer m such that $\gcd(m, (3(n-1))!) = 1$, there exist infinitely many balanced n -simplices of $\mathbb{Z}/m\mathbb{Z}$ generated by PCA_{n-1} , for all possible orientations $\varepsilon \in \{-1, 1\}^n$. In the special case of the two orientations $\varepsilon = + \dots + -$ or $\varepsilon = - \dots - +$, the existence of an infinite number of such balanced simplices is verified for every m such that $\gcd(m, n!) = 1$ for n even and for every m such that $\gcd(m, (\frac{3n+1}{2})!) = 1$ for n odd.*

This paper is organized as follows. After giving some basic results on balanced simplices and orbits of arithmetic arrays generated by ACA in Section 2, we study, in Section 3, arithmetic simplices and we give some sufficient conditions on them to be balanced, for any dimension $n \geq 2$. Moreover, in dimension 2 and 3, we complete by providing necessary conditions on arithmetic triangles and arithmetic tetrahedra for being balanced. This leads to the proof of Theorem 1.14, the main result of this paper, in Section 4. Moreover, using the specificities on balanced arithmetic tetrahedra in dimension 2, highlighted in Section 3, we complete Theorem 1.14 for balanced tetrahedra. In Section 5, we consider the special case where simplices have the additional geometric property of being constituted by antisymmetric sequences. This permits us to obtain more results for ACA of dimension 1 generating balanced triangles. Finally, the new results obtained on balanced simplices generated by PCA_{n-1} are summarized and the problem of determining the existence of balanced ones for the remaining open cases is posed in the last section.

2. PRELIMINARIES

We begin this section with the terminology on simplices that we will use in the sequel.

Definition 2.1 (Vertices, edges, facets and rows of simplices). Let $A = (a_i)_{i \in \mathbb{Z}^n}$ be an infinite array of dimension n of $\mathbb{Z}/m\mathbb{Z}$. Let $\Delta = \Delta(j, \varepsilon, s)$ be the n -simplex of size s of $\mathbb{Z}/m\mathbb{Z}$ with principal vertex at position $j \in \mathbb{Z}^n$ in A and with orientation $\varepsilon \in \{-1, 1\}^n$. Let (e_1, e_2, \dots, e_n) denote the canonical basis of the vector space \mathbb{Z}^n and let $e_0 := 0_{\mathbb{Z}^n}$. The $n + 1$ vertices V_0, \dots, V_n of Δ are defined by $V_k(\Delta) := a_{j+(s-1)\varepsilon \cdot e_k}$ for all $k \in [0, n]$, that are $V_0(\Delta) := a_j$ (principal vertex) and

$$V_k(\Delta) := a_{j+\varepsilon \cdot (s-1)e_k} = a_{j_1, \dots, j_{k-1}, j_k+\varepsilon_k(s-1), j_{k+1}, \dots, j_n},$$

for all $k \in [1, n]$. The $\binom{n+1}{2}$ edges $E_{k,l}$ of Δ are sequences of length s defined by

$$\begin{aligned} E_{k,l}(\Delta) &:= \{a_{j+\varepsilon \cdot ((s-1-x)e_k + xe_l)} \mid x \in [0, s-1]\} \\ &= \{a_{j+\varepsilon \cdot (s-1)e_k}, a_{j+\varepsilon \cdot ((s-2)e_k + e_l)}, a_{j+\varepsilon \cdot ((s-3)e_k + 2e_l)}, \dots, a_{j+\varepsilon \cdot (s-1)e_l}\}, \end{aligned}$$

for all distinct integers $k, l \in [0, n]$. The $n + 1$ facets F_0, \dots, F_n of Δ are the $(n - 1)$ -simplices of size s defined by

$$F_k(\Delta) := \{a_{j+\varepsilon \cdot l} \mid l \in \mathbb{N}^n \text{ such that } l_k = 0 \text{ and } l_1 + \dots + l_n \leq s - 1\},$$

for all $l \in [1, n]$ and

$$F_0(\Delta) := \{a_{j+\varepsilon \cdot l} \mid l \in \mathbb{N}^n \text{ such that } l_1 + \dots + l_n = s - 1\}.$$

For every $k \in [0, s - 1]$, the k th row of Δ is the $(n - 1)$ -simplex of size $s - k$ defined by

$$R_k(\Delta) := \{a_{j+\varepsilon \cdot l} \mid l \in \mathbb{N}^n \text{ such that } l_n = k \text{ and } l_1 + \dots + l_{n-1} \leq s - k - 1\}.$$

2.1. Sizes of balanced simplices. In this subsection, the admissible sizes of balanced simplices are studied. First, the cardinality of an n -simplex of size s is determined.

Proposition 2.2. *Let Δ be an n -simplex of size s appearing in an n -dimensional array. Then, the multiset cardinality of Δ is $|\Delta| = \binom{s+n-1}{n}$.*

Proof. By induction on n . For $n = 1$, Δ is a finite sequence of length s . Thus, $|\Delta| = s = \binom{s}{1}$. Suppose that every $(n - 1)$ -simplex of size k has cardinality of $\binom{k+n-2}{n-1}$, for all integers $k \geq 1$. Now, let Δ be an n -simplex of size s . Since, for all $k \in [0, s - 1]$, the k th row R_k of Δ is an $(n - 1)$ -simplex of size $s - k$, it follows that

$$|\Delta| = \sum_{k=0}^{s-1} |R_k| = \sum_{k=0}^{s-1} \binom{s-k+n-2}{n-1} = \sum_{k=n-1}^{s+n-2} \binom{k}{n-1} = \binom{s+n-1}{n}.$$

This completes the proof. \square

The divisibility of $\binom{s+n-1}{n}$ by m is obviously a necessary condition for having a balanced n -simplex of $\mathbb{Z}/m\mathbb{Z}$ of size s . When m is a composite number, to give all the sizes s for which the binomial $\binom{s+n-1}{n}$ is divisible by m is tedious and not really important here because the results that we obtain in this paper are only for some of them, not for all the admissible sizes. Nevertheless, we can see that the sizes involved in Theorem 1.14 are admissible for this problem.

Proposition 2.3. *Let n, p, k, s be positive integers such that p is prime and $p > n \geq 2$. Then, the binomial coefficient $\binom{s+n-1}{n}$ is divisible by p^k if and only if $s \equiv -t \pmod{p^k}$ for $t \in [0, n - 1]$.*

Proof. Since p^k and $n!$ are relatively prime, we have

$$\binom{s+n-1}{n} \equiv 0 \pmod{p^k} \iff (s+n-1)(s+n-2) \cdots s \equiv 0 \pmod{p^k}.$$

Moreover, $n < p$ implies that p can divide at most one factor of $(s+n-1)(s+n-2)\cdots s$. Therefore, $\binom{s+n-1}{n}$ is divisible by p^k if and only if $s+t \equiv 0 \pmod{p^k}$ for some $t \in [0, n-1]$. This concludes the proof. \square

Proposition 2.4. *Let m and n be two positive integers such that $\gcd(m, n!) = 1$ and let s be a positive integer such that $s \equiv -t \pmod{m}$, where $t \in [0, n-1]$. Then, the binomial coefficient $\binom{s+n-1}{n}$ is divisible by m .*

Proof. If $s \equiv -t \pmod{m}$, then $(s+n-1)(s+n-2)\cdots(s+1)s$ is divisible by m . Finally, since $\gcd(m, n!) = 1$, we deduce from Gauss Lemma that m divides $\frac{(s+n-1)(s+n-2)\cdots(s+1)s}{n!} = \binom{s+n-1}{n}$. This concludes the proof. \square

2.2. Projection Theorem. The result presented in this subsection is useful for proving, by induction on m , that multisets of $\mathbb{Z}/m\mathbb{Z}$ are balanced.

For any divisor α of the positive integer m , denote by π_α the canonical projective map $\pi_\alpha : \mathbb{Z}/m\mathbb{Z} \rightarrow \mathbb{Z}/\alpha\mathbb{Z}$. For any multiset M of $\mathbb{Z}/m\mathbb{Z}$, its projection $\pi_\alpha(M)$ into $\mathbb{Z}/\alpha\mathbb{Z}$ is the multiset of $\mathbb{Z}/\alpha\mathbb{Z}$ defined by $\mathbf{m}_{\pi_\alpha(M)}(y) = \sum_{x \in \pi_\alpha^{-1}(y)} \mathbf{m}_M(x)$ for all $y \in \mathbb{Z}/\alpha\mathbb{Z}$.

Theorem 2.5. *Let m and α be two positive integers such that m is divisible by α and let M be a finite multiset of elements in $\mathbb{Z}/m\mathbb{Z}$. Then, the multiset M is balanced in $\mathbb{Z}/m\mathbb{Z}$ if and only if its projection $\pi_\alpha(M)$ is balanced in $\mathbb{Z}/\alpha\mathbb{Z}$ and $\mathbf{m}_M(x+\alpha) = \mathbf{m}_M(x)$ for all $x \in \mathbb{Z}/m\mathbb{Z}$.*

Proof. Since $\pi_\alpha^{-1}(\{\pi_\alpha(x)\}) = \{x + \lambda\alpha \mid \lambda \in [0, \frac{m}{\alpha} - 1]\}$, for all $x \in \mathbb{Z}/m\mathbb{Z}$, it follows that $\mathbf{m}_{\pi_\alpha(M)}(\pi_\alpha(x)) = \sum_{\lambda=0}^{\frac{m}{\alpha}-1} \mathbf{m}_M(x + \lambda\alpha)$ and the result follows. \square

2.3. Orbits of arithmetic arrays. In this subsection, the orbits of arithmetic arrays are studied in detail. Let $n \geq 2$ be a positive integer. First, we show that the arithmetic structure is preserved under the action of ∂ for any weight array $W = (w_i)_{i \in [-r, r]^{n-1}}$, of radius $r \in \mathbb{N}$.

Proposition 2.6. *Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$. Then,*

$$\partial \text{AA}(a, d) = \text{AA} \left(\sigma a + \sum_{k=1}^{n-1} \sigma_k d_k, \sigma d \right),$$

where σ and σ_k are the coefficients

$$\sigma := \sum_{j \in [-r, r]^{n-1}} w_j, \quad \sigma_k := \sum_{j \in [-r, r]^{n-1}} j_k w_j, \quad \text{for all } k \in [1, n-1].$$

Proof. Let $\text{AA}(a, d) = (a_i)_{i \in \mathbb{Z}^{n-1}}$ and $\partial \text{AA}(a, d) = (b_i)_{i \in \mathbb{Z}^{n-1}}$. By definition of ∂ , for every $i \in \mathbb{Z}^{n-1}$, we have

$$b_i = \sum_{j \in [-r, r]^{n-1}} w_j a_{i+j} = \sum_{j \in [-r, r]^{n-1}} w_j \left(a + \sum_{k=1}^{n-1} (i_k + j_k) d_k \right) = \left(\sigma a + \sum_{k=1}^{n-1} \sigma_k d_k \right) + \sum_{k=1}^{n-1} i_k (\sigma d_k).$$

The result follows. \square

Proposition 2.7. *Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$. Then,*

$$\partial^i \text{AA}(a, d) = \text{AA} \left(\sigma^i a + i \sigma^{i-1} \sum_{k=1}^{n-1} \sigma_k d_k, \sigma^i d \right),$$

for all $i \in \mathbb{N}$.

Proof. By induction on $i \in \mathbb{N}$. For $i = 0$, we retrieve that $\partial^0 \text{AA}(a, d) = \text{AA}(a, d)$. For $i \geq 1$, by the recursive definition of ∂^i and Proposition 2.6, we obtain that

$$\begin{aligned} \partial^i \text{AA}(a, d) &= \partial (\partial^{i-1} \text{AA}(a, d)) \\ &= \partial \left(\text{AA} \left(\sigma^{i-1} a + (i-1) \sigma^{i-2} \sum_{k=1}^{n-1} \sigma_k d_k, \sigma^{i-1} d \right) \right) \\ &= \text{AA} \left(\sigma \left(\sigma^{i-1} a + (i-1) \sigma^{i-2} \sum_{k=1}^{n-1} \sigma_k d_k \right) + \sum_{k=1}^{n-1} \sigma_k (\sigma^{i-1} d_k), \sigma (\sigma^{i-1} d) \right) \\ &= \text{AA} \left(\sigma^i a + i \sigma^{i-1} \sum_{k=1}^{n-1} \sigma_k d_k, \sigma^i d \right). \end{aligned}$$

This concludes the proof. \square

Thus, the elements of the orbit of an arithmetic array $\text{AA}(a, d)$ are entirely determined in function of a, d, σ and σ_k for all $k \in [1, n-1]$.

Proposition 2.8. *Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$. Let $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^{n-1} \times \mathbb{N}}$ be the orbit of the arithmetic array $\text{AA}(a, d)$. Then,*

$$a_i = (\partial^{i_n} \text{AA}(a, d))_{i_1, \dots, i_{n-1}} = \sigma^{i_n} \left(a + \sum_{k=1}^{n-1} i_k d_k \right),$$

for all $i \in \mathbb{Z}^{n-1} \times \mathbb{N}$, where $d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k$.

Proof. Directly comes from Proposition 2.7. \square

Remark 2.9. For $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^{n-1} \times \mathbb{N}}$ and for every $(i_1, \dots, i_{n-1}) \in \mathbb{Z}^{n-1}$, the sequence $(a_{i_1, \dots, i_{n-1}, i_n})_{i_n \in \mathbb{N}}$ is the arithmetico-geometric sequence with first element $a + i_1 d_1 + \dots + i_{n-1} d_{n-1}$, with common difference $d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k$ and common ratio σ .

We deduce from Proposition 2.8 that two distinct ACA can generate the same orbit from an arithmetic array. For instance, for any ACA of weight array $W = (w_i)_{i \in [-r, r]^{n-1}}$ of radius r , we can consider the ACA of weight array $\overline{W} = (\overline{w}_i)_{i \in [-1, 1]^{n-1}}$ of radius 1 defined by

$$\overline{w}_i = \begin{cases} \sigma(W) - \sum_{k=1}^{n-1} \sigma_k(W) & , \text{ if } i = 0_{\mathbb{Z}^n}, \\ \sigma_k(W) & , \text{ if } i = e_k, \text{ for all } k \in [1, n-1], \\ 0 & , \text{ otherwise.} \end{cases}$$

Then, it is clear that we have

$$\sigma(\overline{W}) = \sum_{i \in [-1, 1]^{n-1}} \overline{w}_i = \sigma(W),$$

and

$$\sigma_k(\overline{W}) = \sum_{i \in [-1, 1]^{n-1}} i_k \overline{w}_i = \sigma_k(W),$$

for all $k \in [1, n-1]$. Therefore, in the sequel of this paper, the coefficients σ and σ_k will be more important than the elements of the weight array W themselves.

Now, we prove that, in the orbit of an arithmetic array of $\mathbb{Z}/m\mathbb{Z}$, if there exists a balanced simplex of sufficiently large size, then σ is invertible modulo m .

Proposition 2.10. *Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$. In the orbit $\mathcal{O}(\text{AA}(a, d))$, if an n -simplex of size $s \geq \lceil \frac{5n+3}{2} \rceil$ is balanced, then σ is invertible modulo m .*

Proof. Let $j \in \mathbb{Z}^{n-1} \times \mathbb{N}$ and let $\varepsilon \in \{-1, +1\}^n$. Suppose that $\Delta(j, \varepsilon, s)$ is balanced and that σ is not invertible modulo m . Without loss of generality, we can suppose that $\sigma \equiv 0 \pmod{m}$. If not, we consider the projection into $\mathbb{Z}/\text{gcd}(\sigma, m)\mathbb{Z}$ and then we have $\sigma \equiv 0 \pmod{\text{gcd}(\sigma, m)}$. By Proposition 2.7, we know that, in the case $\sigma \equiv 0 \pmod{m}$, we have $\partial^i \text{AA}(a, d) = \text{AA}(0, 0)$, the constant array equal to zero in $\mathbb{Z}/m\mathbb{Z}$, for all $i \geq 2$. Therefore, all the elements in the k th row of $\Delta(j, \varepsilon, s)$ are constituted by elements equal to zero, for all k such that $j_n + k\varepsilon_n \geq 2$. Thus $\Delta(j, \varepsilon, s)$ contains at least $\binom{s+n-3}{n}$ elements equal to zero by Proposition 2.2. Moreover, since $\Delta(j, \varepsilon, s)$ is balanced, we deduce that $\binom{s+n-3}{n} \leq \binom{s+n-1}{n} - \binom{s+n-3}{n}$ since $\Delta(j, \varepsilon, s)$ must contain all the other elements of $\mathbb{Z}/m\mathbb{Z}$ with the same multiplicity. It follows that

$$\binom{s+n-3}{n} \leq \binom{s+n-2}{n-1} + \binom{s+n-3}{n-1}.$$

Since

$$\begin{aligned} & \binom{s+n-3}{n} + \binom{s+n-2}{n-1} + \binom{s+n-3}{n-1} \\ &= \binom{s+n-3}{n} + 2\binom{s+n-3}{n-1} + \binom{s+n-3}{n-2} = \binom{s+n-1}{n}, \end{aligned}$$

it follows that we have

$$\begin{aligned} 2\binom{s+n-3}{n} \leq \binom{s+n-1}{n} &\iff 2 \prod_{i=s-2}^{s+n-3} i \leq \prod_{i=s}^{s+n-1} i \\ &\iff 2(s-2)(s-1) \leq (s+n-2)(s+n-1). \end{aligned}$$

Thus $s^2 - (2n+3)s - (n^2 - 3n - 2) \leq 0$ and we deduce that it is possible only if

$$s \leq \frac{2n+3 + \sqrt{8n^2+1}}{2} < \frac{5n+3}{2},$$

in contradiction with the hypothesis that $s \geq \lceil \frac{5n+3}{2} \rceil$. This concludes the proof. \square

This is the reason why we suppose, in the sequel of this paper, that σ is invertible modulo m . We end this section by showing that a simplex in the orbit of an arithmetic array can be decomposed into arithmetic subsimplices.

Proposition 2.11. *Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$. Let α and s be two positive integers such that α is divisible by $\text{ord}_m(\sigma)$ and $s \equiv -t \pmod{\alpha}$, where $t \in [0, n-1]$, and let $\varepsilon \in \{-1, 1\}^n$. Let $\Delta(j, \varepsilon, s)$ be the n -simplex appearing in the orbit $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^{n-1} \times \mathbb{N}}$. Then, for every $k \in [0, \alpha-1]^n$, the subsimplex*

$$\text{SS}_k := \left\{ a_{j+\varepsilon \cdot (k+\alpha l)} \mid l \in \mathbb{N}^n \text{ such that } (k_1 + \alpha l_1) + \dots + (k_n + \alpha l_n) \leq s-1 \right\},$$

obtained from $\Delta(j, \varepsilon, s)$ by extracting one term every α in each component, is the arithmetic simplex

$$\text{SS}_k = \text{AS} \left(a_{j+\varepsilon \cdot k}, \alpha \sigma^{j_n + \varepsilon_n k_n} \varepsilon \cdot \tilde{d}, \left\lceil \frac{s}{\alpha} \right\rceil - \left\lfloor \frac{\sum_{u=1}^n k_u + t}{\alpha} \right\rfloor \right),$$

where $\tilde{d} = (d_1, \dots, d_{n-1}, \sigma^{-1} \sum_{u=1}^{n-1} \sigma_u d_u)$.

Proof. Let $k \in [0, \alpha - 1]^n$. As already observed in Remark 2.9, for every $(i_1, \dots, i_{n-1}) \in \mathbb{Z}^{n-1}$, the sequence $(a_{i_1, \dots, i_{n-1}, i_n})_{i_n \in \mathbb{N}}$ is the arithmetico-geometric sequence whose first element is $a + i_1 d_1 + \dots + i_{n-1} d_{n-1}$, with common difference $\sigma^{-1} \sum_{u=1}^{n-1} \sigma_u d_u$ and with common ratio σ . Since α is a multiple of $\text{ord}_m(\sigma)$, it follows that, for every $(l_1, \dots, l_{n-1}) \in \mathbb{N}^{n-1}$, the sequence $(a_{j+\varepsilon \cdot (k+\alpha l)})_{l_n \in \mathbb{N}}$ is arithmetic, with common difference $\alpha \sigma^{j_n + \varepsilon_n k_n - 1} \varepsilon_n \sum_{u=1}^{n-1} \sigma_u d_u$. Indeed, from Proposition 2.8, we obtain

$$a_{j+\varepsilon \cdot (k+\alpha l)} = \sigma^{j_n + \varepsilon_n (k_n + \alpha l_n)} \left(a + \sum_{u=1}^n (j_u + \varepsilon_u (k_u + \alpha l_u)) d_u \right),$$

where $d_n := \sigma^{-1} \sum_{u=1}^{n-1} \sigma_u d_u$. Since α is a multiple of $\text{ord}_n(\sigma)$, we have

$$\sigma^{j_n + \varepsilon_n (k_n + \alpha l_n)} = \sigma^{j_n + \varepsilon_n k_n}.$$

Thus

$$a_{j+\varepsilon \cdot (k+\alpha l)} = \sigma^{j_n + \varepsilon_n k_n} \left[\left(a + \sum_{u=1}^n (j_u + \varepsilon_u k_u) d_u \right) + \alpha \sum_{u=1}^n l_u \varepsilon_u d_u \right].$$

By Proposition 2.8 again, for $a_{j+\varepsilon \cdot k}$,

$$a_{j+\varepsilon \cdot (k+\alpha l)} = a_{j+\varepsilon \cdot k} + \alpha \sigma^{j_n + \varepsilon_n k_n} \sum_{u=1}^n l_u (\varepsilon_u d_u),$$

for every $l \in \mathbb{N}^n$. Therefore the subsimplex SS_k is an arithmetic simplex whose principal vertex is $a_{j+\varepsilon \cdot k}$ and with common difference $\alpha \sigma^{j_n + \varepsilon_n k_n} \varepsilon \cdot \tilde{d}$, where $\tilde{d} = (d_1, d_2, \dots, d_n)$. It remains to determine the size of this arithmetic simplex. Let $\lambda := \lceil \frac{s}{\alpha} \rceil$ and $\mu := \lfloor \frac{\sum_{u=1}^n k_u + t}{\alpha} \rfloor$. In other words, we have $s = \lambda \alpha - t$ and

$$\mu \alpha - t \leq \sum_{u=1}^n k_u \leq (\mu + 1) \alpha - t - 1.$$

For any $l \in \mathbb{N}^n$, the inequality

$$\sum_{u=1}^n (k_u + \alpha l_u) \leq s - 1 = \lambda \alpha - t - 1$$

is then equivalent to

$$\sum_{u=1}^n l_u \leq \lambda - \mu - 1.$$

Therefore the arithmetic simplex SS_k is of size $\lambda - \mu$. This completes the proof. \square

From the previous proposition, we know that every n -simplex Δ of size $\lambda \alpha - t$, where α is a multiple of $\text{ord}_m(\sigma)$ and $t \in [0, n - 1]$, appearing in the orbit of an arithmetic array can be decomposed into α^n arithmetic n -simplices of sizes in $[\lambda - (n - 1), \lambda]$. Therefore, in next section, the arithmetic simplices will be studied in detail.

3. BALANCED ARITHMETIC SIMPLICES

In this section, we will see that arithmetic simplices are a source of balanced multisets of $\mathbb{Z}/m\mathbb{Z}$. First, we show, in the general case $n \geq 1$, that there exists sufficient conditions on arithmetic simplices for being balanced. After that, in dimension $n = 2$ and $n = 3$, i.e. for arithmetic triangles and arithmetic tetrahedra, necessary conditions for being balanced are also given.

3.1. The general case : in dimension $n \geq 1$. We begin this subsection by showing that, when $n \geq 2$, the edges, the facets and the rows of an arithmetic simplex are also arithmetic.

Proposition 3.1. *Let $a \in \mathbb{Z}/m\mathbb{Z}$, $d = (d_1, \dots, d_n) \in (\mathbb{Z}/m\mathbb{Z})^n$ and let s be a positive integer. Let $\Delta := \text{AS}(a, d, s)$ and let $d_0 := 0$. Then, we have*

$$V_i(\Delta) = a + (s - 1)d_i,$$

for all $i \in [0, n]$,

$$E_{i,j}(\Delta) = \text{AP}(V_i(\Delta), d_j - d_i, s),$$

for all distinct integers $i, j \in [0, n]$,

$$R_i(\Delta) = \text{AS}(a + id_n, (d_1, \dots, d_{n-1}), s - i),$$

for all $i \in [0, n]$,

$$F_i(\Delta) = \text{AS}(a, (d_1, \dots, d_{i-1}, d_{i+1}, \dots, d_n), s),$$

for all $i \in [1, n]$, and

$$F_0(\Delta) = \text{AS}(a + (s - 1)d_1, (d_2 - d_1, \dots, d_n - d_1), s).$$

Moreover, for all $i \in [0, n]$, we have

$$\Delta \setminus F_i(\Delta) = \text{AS}(a + d_i, d, s - 1).$$

Proof. By Definition 1.12 and Definition 2.1. □

The following theorem, which gives sufficient conditions on arithmetic simplices for being balanced, is the main result of this section.

Theorem 3.2. *Let n and m be two positive integers such that $\gcd(m, n!) = 1$. Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d = (d_1, \dots, d_n) \in (\mathbb{Z}/m\mathbb{Z})^n$ such that d_i , for all $1 \leq i \leq n$, and $d_j - d_i$, for all $1 \leq i < j \leq n$, are invertible. Then, the arithmetic simplex $\text{AS}(a, d, s)$ is balanced for all $s \equiv -t \pmod{m}$, with $t \in [0, n - 1]$.*

The proof of this theorem is based on the following result, which is a key lemma in this paper.

Lemma 3.3. *Let $n \geq 2$, m and s be positive integers. Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d \in (\mathbb{Z}/m\mathbb{Z})^n$. Then, the multiplicity function of $\Delta = \text{AS}(a, d, s)$ verifies*

$$\mathbf{m}_\Delta(x + d_j) - \mathbf{m}_\Delta(x + d_i) = \mathbf{m}_{F_j(\Delta)}(x + d_j) - \mathbf{m}_{F_i(\Delta)}(x + d_i),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$ and for all distinct integers $i, j \in [0, n]$, where $d_0 := 0$.

Proof. Let $0 \leq i < j \leq n$. Since, from Proposition 3.1, we have

$$\Delta \setminus F_k(\Delta) = \text{AS}(a + d_k, d, s - 1),$$

for all integers $k \in [0, n]$, it follows that the arithmetic simplex $\Delta \setminus F_j(\Delta)$ is a translate of $\Delta \setminus F_i(\Delta)$ by $d_j - d_i$. Thus, for all $x \in \mathbb{Z}/m\mathbb{Z}$, we have

$$\mathbf{m}_{\Delta \setminus F_j(\Delta)}(x + d_j) = \mathbf{m}_{\Delta \setminus F_i(\Delta)}(x + d_i).$$

Therefore,

$$\mathbf{m}_\Delta(x + d_j) - \mathbf{m}_{F_j(\Delta)}(x + d_j) = \mathbf{m}_\Delta(x + d_i) - \mathbf{m}_{F_i(\Delta)}(x + d_i)$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. □




FIGURE 4. Common differences of arithmetic triangles

Proof of Theorem 3.2. By induction on n .

For $n = 1$ and $s \equiv 0 \pmod{m}$, the arithmetic simplex $\text{AS}(a, d, s)$ is simply an arithmetic progression with invertible common difference $d \in \mathbb{Z}/m\mathbb{Z}$ and of length a multiple of m . Therefore $\text{AS}(a, d, s)$ is balanced in this case.

Suppose now that $n \geq 2$ and that the result is true in dimension $\leq n - 1$. We distinguish different cases depending on the residue class of s modulo m .

Case 1. For $s \equiv -t \pmod{m}$, with $t \in [0, n - 2]$.

Let $\Delta = \text{AS}(a, d, s)$. First, from Proposition 3.1, we know that

$$F_1(\Delta) = \text{AS}(a, (d_2, \dots, d_n), s) \text{ and } F_2(\Delta) = \text{AS}(a, (d_1, d_3, \dots, d_n), s).$$

Moreover, since d_i , for all $1 \leq i \leq n$, and $d_j - d_i$, for all $1 \leq i < j \leq n$, are invertible, we obtain by induction hypothesis that the facets $F_1(\Delta)$ and $F_2(\Delta)$ are balanced simplices of dimension $n - 1$. Therefore, their multiplicity functions $\mathbf{m}_{F_1(\Delta)}$ and $\mathbf{m}_{F_2(\Delta)}$ are constant on $\mathbb{Z}/m\mathbb{Z}$, equal to

$$\mathbf{m}_{F_1(\Delta)}(x) = \mathbf{m}_{F_2(\Delta)}(x) = \frac{1}{m} \binom{s + n - 2}{n - 1},$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. By Lemma 3.3, we obtain

$$\mathbf{m}_\Delta(x + d_1) - \mathbf{m}_\Delta(x + d_2) = \mathbf{m}_{F_1(\Delta)}(x + d_1) - \mathbf{m}_{F_2(\Delta)}(x + d_2) = 0,$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Finally, since $d_2 - d_1$ is invertible and

$$\mathbf{m}_\Delta(x + (d_1 - d_2)) = \mathbf{m}_\Delta(x)$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$, we conclude that \mathbf{m}_Δ is a constant function and so the arithmetic simplex Δ is balanced in $\mathbb{Z}/m\mathbb{Z}$.

Case 2. For $s \equiv -(n - 1) \pmod{m}$.

The arithmetic simplex $\text{AS}(a, (d_1, \dots, d_n), s)$ can be seen as the multiset difference $\Delta \setminus F_1(\Delta)$, where

$$\Delta = \text{AS}(a - d_1, (d_1, \dots, d_n), s + 1).$$

Since $s + 1 \equiv -(n - 2) \pmod{m}$, it follows, from Case 1 and from the induction hypothesis, respectively, that $\Delta = \text{AS}(a - d_1, (d_1, \dots, d_n), s + 1)$ and $F_1(\Delta) = \text{AS}(a - d_1, (d_2, \dots, d_n), s + 1)$ are balanced. The multiset difference of balanced multisets is also balanced. This concludes the proof. \square

3.2. In dimension 2. In this subsection, we only consider arithmetic triangles over $\mathbb{Z}/m\mathbb{Z}$. Necessary conditions on the common differences d_1 , d_2 and $d_2 - d_1$ of $\text{AS}(a, (d_1, d_2), s)$, depicted in Figure 4, for being balanced in $\mathbb{Z}/m\mathbb{Z}$ are determined.

Theorem 3.4. *Let m and s be two positive integers and let $a, d_1, d_2 \in \mathbb{Z}/m\mathbb{Z}$. If the arithmetic triangle $\text{AS}(a, (d_1, d_2), s)$ is balanced, then the common differences d_1 , d_2 and $d_2 - d_1$ are all invertible.*

The proof of this theorem is based on the following lemma, where the multiplicity function of the arithmetic triangle $\text{AS}(a, (0, d), s)$, with d invertible, is explicitly given.

Lemma 3.5. *Let m and s be two positive integers and let $a, d \in \mathbb{Z}/m\mathbb{Z}$ such that d is invertible. Then, the arithmetic triangle $\Delta = \text{AS}(a, (0, d), s)$ is not balanced in $\mathbb{Z}/m\mathbb{Z}$. Moreover, if $s = \lambda m + \mu$ is the Euclidean division of s by m , we have*

$$\mathbf{m}_\Delta(a + id) = \binom{\lambda + 1}{2} m + \left\lceil \frac{s - i}{m} \right\rceil (\mu - i),$$

for all integers $i \in [0, m - 1]$.

Proof. Let $s = \lambda m + \mu$ be the Euclidean division of s by m . As represented in Figure 4, the common differences of $\Delta = \text{AS}(a, (0, d), s)$ are 0 , d and d . Then, for every integer $j \in [0, s - 1]$, the j th row R_j of Δ is the constant sequence of length $s - j$ equal to $a + jd$, that is, $R_j(\Delta) = \text{AP}(a + jd, 0, s - j)$. Thus, the multiplicity function \mathbf{m}_Δ is determined by

$$\begin{aligned} \mathbf{m}_\Delta(a + id) &= \sum_{j=0}^{s-1} \mathbf{m}_{R_j(\Delta)}(a + id) = \sum_{j=0}^{\left\lceil \frac{s-i}{m} \right\rceil} \mathbf{m}_{R_{jm+i}(\Delta)}(a + id) \\ &= \sum_{j=0}^{\left\lceil \frac{s-i}{m} \right\rceil} (s - (jm + i)) = \sum_{j=0}^{\left\lceil \frac{s-i}{m} \right\rceil} ((\lambda - j)m + (\mu - i)) \\ &= \binom{\lambda + 1}{2} m + \left\lceil \frac{s - i}{m} \right\rceil (\mu - i) \end{aligned}$$

for all integers $i \in [0, m - 1]$. It follows that

$$\mathbf{m}_\Delta(a) > \mathbf{m}_\Delta(a + d) > \mathbf{m}_\Delta(a + 2d) > \dots > \mathbf{m}_\Delta(a + (m - 1)d).$$

Since its multiplicity function is not constant on $\mathbb{Z}/m\mathbb{Z}$, the arithmetic triangle Δ is not balanced. This completes the proof. \square

Proof of Theorem 3.4. Let $\Delta = \text{AS}(a, (d_1, d_2), s)$ be an arithmetic triangle and suppose that there exists at least one common difference that is not invertible. Without loss of generality, suppose that d_1 is not invertible. Then, we consider the projection of Δ into $\mathbb{Z}/\alpha\mathbb{Z}$, where $\alpha = \gcd(d_1, m) \geq 2$. Then,

$$\pi_\alpha(\Delta) = \text{AS}(\pi_\alpha(a), (0, \pi_\alpha(d_2)), s).$$

If $\pi_\alpha(d_2)$ is invertible in $\mathbb{Z}/\alpha\mathbb{Z}$, then Δ is not balanced in $\mathbb{Z}/m\mathbb{Z}$ since $\pi_\alpha(\Delta)$ is not balanced in $\mathbb{Z}/\alpha\mathbb{Z}$ by Lemma 3.5. Otherwise, if $\pi_\alpha(d_2)$ is not invertible in $\mathbb{Z}/\alpha\mathbb{Z}$, the result follows since the projection of $\pi_\alpha(\Delta)$ into $\mathbb{Z}/\beta\mathbb{Z}$, where $\beta = \gcd(\pi_\alpha(d_1), \alpha) \geq 2$, is the constant triangle uniquely constituted by elements $\pi_\beta(\pi_\alpha(a))$, which is obviously not balanced in $\mathbb{Z}/\beta\mathbb{Z}$. \square

It follows from this theorem that there does not exist balanced arithmetic triangles in $\mathbb{Z}/m\mathbb{Z}$ for m even. Nevertheless, in the case where m is an even number, the multiplicity function of an arithmetic triangle of $\mathbb{Z}/m\mathbb{Z}$ can be completely determined when exactly two of the three common differences $d_1, d_2, d_1 - d_2$ are invertible and the size s is such that $s \equiv 0$ or $-1 \pmod{m}$.

Proposition 3.6. *Let m and s be two positive integers such that $s \equiv 0$ or $-1 \pmod{m}$. Let $a, d_1, d_2 \in \mathbb{Z}/m\mathbb{Z}$ and let $\Delta = \text{AS}(a, (d_1, d_2), s)$. If d_2 and $d_2 - d_1$ are invertible, then*

$$\mathbf{m}_\Delta(x) = \mathbf{m}_\Delta(x + \gcd(d_1, m)),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$, and

$$\mathbf{m}_\Delta(a + id_2) = \frac{1}{m} \binom{s+1}{2} + \left\lceil \frac{s}{m} \right\rceil \left(\frac{\gcd(d_1, m) - 1}{2} - i \right),$$

for all integers $i \in [0, \gcd(d_1, m) - 1]$.

Proof. Since d_2 and $d_2 - d_1$ are invertible and $s \equiv 0$ or $-1 \pmod{m}$, it follows that $F_1(\Delta) = \text{AP}(a, d_2, s)$ and $F_0(\Delta) = \text{AP}(a + (s-1)d_1, d_2 - d_1, s)$ are balanced in $\mathbb{Z}/m\mathbb{Z}$. Therefore, $\mathbf{m}_{F_1(\Delta)}(x) = \mathbf{m}_{F_0(\Delta)}(x) = \frac{1}{m} \binom{s+1}{2}$ for all $x \in \mathbb{Z}/m\mathbb{Z}$. It follows, from Lemma 3.3, that

$$\mathbf{m}_\Delta(x) - \mathbf{m}_\Delta(x + d_1) = \mathbf{m}_{F_0(\Delta)}(x) - \mathbf{m}_{F_1(\Delta)}(x + d_1) = 0,$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. The first identity is then proved.

The second identity, in the case where d_1 is invertible, comes from the fact that Δ is balanced, by Theorem 3.2, in this case. Suppose now that $\alpha = \gcd(d_1, m) \geq 2$ and consider the projected triangle $\pi_\alpha(\Delta)$ in $\mathbb{Z}/\alpha\mathbb{Z}$. Then, since $\pi_\alpha(\Delta) = \text{AS}(\pi_\alpha(a), (0, \pi_\alpha(d_2)), s)$ and $\pi_\alpha(d_2)$ is invertible in $\mathbb{Z}/\alpha\mathbb{Z}$, the multiplicity function $\mathbf{m}_{\pi_\alpha(\Delta)}$ is entirely determined by Lemma 3.5. Let $s = \lambda m + \mu$ be the Euclidean division of s by m , where $\mu \in \{0, m-1\}$. Then, the Euclidean division of s by α is $s = \lambda \frac{m}{\alpha} \alpha$, if $\mu = 0$, and $((\lambda + 1) \frac{m}{\alpha} - 1) \alpha + (\alpha - 1)$, if $\mu = m - 1$. The multiplicity function $\mathbf{m}_{\pi_\alpha(\Delta)}$ verifies

$$\mathbf{m}_{\pi_\alpha(\Delta)}(\pi_\alpha(a) + i\pi_\alpha(d)) = \begin{cases} \left(\lambda \frac{m}{\alpha} + 1 \right) \alpha - \left\lceil \frac{s-i}{\alpha} \right\rceil i & , \text{ if } \mu = 0, \\ \left((\lambda + 1) \frac{m}{\alpha} \right) \alpha + \left\lceil \frac{s-i}{\alpha} \right\rceil (\alpha - 1 - i) & , \text{ if } \mu = m - 1, \end{cases}$$

for all $i \in [0, \alpha - 1]$. Moreover, by the first identity, since $\mathbf{m}_\Delta(x + \alpha) = \mathbf{m}_\Delta(x)$ for all $x \in \mathbb{Z}/m\mathbb{Z}$, we have

$$\mathbf{m}_\Delta(a + id) = \frac{\alpha}{m} \times \mathbf{m}_{\pi_\alpha(\Delta)}(\pi_\alpha(a) + i\pi_\alpha(d)),$$

for all $i \in [0, \alpha - 1]$. Finally, since

$$\left\lceil \frac{s-i}{\alpha} \right\rceil = \begin{cases} \lambda \frac{m}{\alpha} & , \text{ if } \mu = 0, i \in [0, \alpha - 1], \\ (\lambda + 1) \frac{m}{\alpha} & , \text{ if } \mu = m - 1, i \in [0, \alpha - 2], \\ (\lambda + 1) \frac{m}{\alpha} - 1 & , \text{ if } \mu = m - 1, i = \alpha - 1, \end{cases}$$

it follows that

$$\mathbf{m}_\Delta(a + id) = \left(\lambda \frac{m}{\alpha} + 1 \right) \frac{\alpha^2}{m} - \lambda i = \frac{\lambda(\lambda m + \alpha)}{2} - \lambda i = \frac{1}{m} \binom{\lambda m + 1}{2} + \lambda \left(\frac{\alpha - 1}{2} - i \right),$$

for all $i \in [0, \alpha - 1]$ if $\mu = 0$, and

$$\begin{aligned} \mathbf{m}_\Delta(a + id) &= \left((\lambda + 1) \frac{m}{\alpha} \right) \frac{\alpha^2}{m} + (\lambda + 1)(\alpha - 1 - i) \\ &= \frac{(\lambda + 1)((\lambda + 1)m - \alpha)}{2} + (\lambda + 1)(\alpha - 1 - i) \\ &= \frac{1}{m} \binom{(\lambda + 1)m}{2} + (\lambda + 1) \left(\frac{\alpha - 1}{2} - i \right), \end{aligned}$$

x	0	1	2	3	4	5	6	7	8	9	10	11
$\mathbf{m}_\Delta(x)$	5	6	7	8	5	6	7	8	5	6	7	8



TABLE 1. Multiplicity function of $\Delta = \text{AS}(0, (1, 5), 12)$ in $\mathbb{Z}/12\mathbb{Z}$ 

FIGURE 5. Common differences of an arithmetic tetrahedron

FIGURE 6. Non adjacent common differences of $\text{AS}(a, (d_1, d_2, d_3), s)$

for all $i \in [0, \alpha - 1]$ if $\mu = m - 1$. This completes the proof. \square

For example, for $m = 12$, $a = 0$, $d_1 = 1$ and $d_2 = 5$, we obtain that $d_2 - d_1 = 4$, $\gcd(d_1 - d_2, m) = 4$ and the multiplicity function of $\Delta = \text{AS}(a, (d_1, d_2), m)$ is given in Table 1.

3.3. In dimension 3. In this subsection, we only consider the arithmetic tetrahedron $\text{AS}(a, (d_1, d_2, d_3), s)$ in $\mathbb{Z}/m\mathbb{Z}$. We determine necessary and sufficient conditions on the common differences $d_1, d_2, d_3, d_2 - d_1, d_3 - d_2$ and $d_1 - d_3$ of $\text{AS}(a, (d_1, d_2, d_3), s)$, depicted in Figure 5, for being balanced in $\mathbb{Z}/m\mathbb{Z}$.

Definition 3.7 (Adjacent common differences). Among the six common differences $d_1, d_2, d_3, d_2 - d_1, d_3 - d_2$ and $d_1 - d_3$ of $\text{AS}(a, (d_1, d_2, d_3), s)$, two of them are said to be adjacent if they have a vertex in common. The couple of non adjacent common differences of $\text{AS}(a, (d_1, d_2, d_3), s)$ are $(d_1, d_3 - d_2)$, $(d_2, d_1 - d_3)$ and $(d_3, d_2 - d_1)$. The twelve other couples of common differences are said to be adjacent (See Figure 6).

Theorem 3.8. *Let m and s be two positive integers and let $a, d_1, d_2, d_3 \in \mathbb{Z}/m\mathbb{Z}$. Let $D := \{d_1, d_2, d_3, d_2 - d_1, d_3 - d_2, d_1 - d_3\}$ be the set of common differences of the arithmetic tetrahedron $\Delta = \text{AS}(a, (d_1, d_2, d_3), s)$. If Δ is balanced in $\mathbb{Z}/m\mathbb{Z}$ and*

- i) m is odd, then all the elements of D are invertible.*
- ii) m is even, then all the elements of D are invertible, except two of them, say δ_1 and δ_2 , which are non adjacent and such that $\gcd(\delta_1, m) = \gcd(\delta_2, m) = 2$.*




FIGURE 7. Arithmetic tetrahedra with a common difference equal to zero

The proof of this theorem is based on the following four lemmas, where we study the multiplicity function of the arithmetic tetrahedra with at least one common difference equal to zero. A representation of the common differences of these arithmetic tetrahedra can be found in Figure 7 and Figure 8.

Lemma 3.9. *Let m and s be two positive integers and let $a, d \in \mathbb{Z}/m\mathbb{Z}$ such that d is invertible. Then, the arithmetic tetrahedron $AS(a, (0, 0, d), s)$ is not balanced in $\mathbb{Z}/m\mathbb{Z}$.*

Proof. Let $\Delta = AS(a, (0, 0, d), s)$. By Lemma 3.3, we obtain

$$\mathbf{m}_{\Delta}(a) - \mathbf{m}_{\Delta}(a + d) = \mathbf{m}_{F_1(\Delta)}(a) - \mathbf{m}_{F_3(\Delta)}(a + d).$$

Since $F_3(\Delta) = AS(a, (0, 0), s)$, i.e. the constant triangle of size s where all the terms are equal to a , we have $\mathbf{m}_{F_3(\Delta)}(a + d) = 0$. Since $F_1(\Delta) = AS(a, (0, d), s)$, it follows from Lemma 3.5 that

$$\mathbf{m}_{F_1(\Delta)}(a) = \binom{\lambda + 1}{2} m + \left\lceil \frac{s - i}{m} \right\rceil \mu > 0,$$

where $s = \lambda m + \mu$ is the Euclidean division of s by m . This leads to the inequality $\mathbf{m}_{\Delta}(a + d) < \mathbf{m}_{\Delta}(a)$ and thus the multiplicity function \mathbf{m}_{Δ} is not constant on $\mathbb{Z}/m\mathbb{Z}$. This concludes the proof. \square

Lemma 3.10. *Let m and s be two positive integers and let $a, d_1, d_2 \in \mathbb{Z}/m\mathbb{Z}$ such that d_1, d_2 and $d_2 - d_1$ are invertible. Then, the arithmetic tetrahedron $AS(a, (0, d_1, d_2), s)$ is not balanced in $\mathbb{Z}/m\mathbb{Z}$.*

Proof. First, since d_1, d_2 and $d_2 - d_1$ are invertible, we know that m is an odd number. Assume, without loss of generality, that m is an odd prime. Indeed, if m is composite, we know that Δ cannot be balanced in $\mathbb{Z}/m\mathbb{Z}$ if its projected tetrahedron $\pi_p(\Delta)$ is not balanced in $\mathbb{Z}/p\mathbb{Z}$, where p is an odd prime factor of m . Moreover, a necessary condition on s for Δ being balanced is that m divides $\binom{s+2}{3}$. Since m is an odd prime, this implies that $s \equiv 0, -1$ or $-2 \pmod{m}$. If $s \equiv -2 \pmod{m}$, then $\Delta = AS(a, (0, d_1, d_2), s)$ can be seen as $\Delta' \setminus F_1(\Delta')$, where $\Delta' = AS(a, (0, d_1, d_2), s + 1)$ and $F_1(\Delta') = AS(a, (d_1, d_2), s + 1)$. Since $s + 1 \equiv -1 \pmod{m}$ and $d_1, d_2, d_2 - d_1$ are invertible, we deduce from Theorem 3.2 that $F_1(\Delta')$ is balanced. Therefore Δ is balanced if and only if Δ' is. This is the reason why, in this proof, we suppose that m is an odd prime and that s is a positive integer such that $s \equiv 0$ or $-1 \pmod{m}$. First, by Lemma 3.3, we obtain

$$\mathbf{m}_{\Delta}(x + d_2) - \mathbf{m}_{\Delta}(x) = \mathbf{m}_{F_3(\Delta)}(x + d_2) - \mathbf{m}_{F_1(\Delta)}(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Since d_1, d_2 and $d_2 - d_1$ are invertible and $s \equiv 0$ or $-1 \pmod{m}$, we know, from Theorem 3.2, that $F_1(\Delta) = AS(a, (d_1, d_2), s)$ is a balanced triangle. Therefore,

$$\mathbf{m}_{F_1(\Delta)}(x) = \frac{1}{m} \binom{s + 1}{2},$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Moreover, since d_1 is invertible, it follows from Lemma 3.5 that the multiplicity function of $F_3(\Delta) = \text{AS}(a, (0, d_1), s)$ verifies

$$\mathbf{m}_{F_3(\Delta)}(a) > \mathbf{m}_{F_3(\Delta)}(a + d_1) > \cdots > \mathbf{m}_{F_3(\Delta)}(a + (m-1)d_1).$$

Therefore, since $\sum_{i=0}^{m-1} \mathbf{m}_{F_3(\Delta)}(a + id) = \binom{s+1}{2}$, we have $\mathbf{m}_{F_3(\Delta)}(a) > \frac{1}{m} \binom{s+1}{2}$. This leads to the inequality

$$\mathbf{m}_{\Delta}(a) - \mathbf{m}_{\Delta}(a - d_2) = \mathbf{m}_{F_3(\Delta)}(a) - \mathbf{m}_{F_1(\Delta)}(a - d_2) = \mathbf{m}_{F_3(\Delta)}(a) - \frac{1}{m} \binom{s+1}{2} > 0.$$

Thus, the multiplicity function \mathbf{m}_{Δ} is not constant on $\mathbb{Z}/m\mathbb{Z}$. This concludes the proof. \square

Lemma 3.11. *Let m and s be two positive integers and let $a, d \in \mathbb{Z}/m\mathbb{Z}$ such that d is invertible. If the arithmetic tetrahedron $\text{AS}(a, (0, d, d), s)$ is balanced, then $m = 2$ and s is even.*

Proof. Let $\Delta = \text{AS}(a, (0, d, d), s)$. First, by Lemma 3.3, we obtain the identity

$$\mathbf{m}_{\Delta}(x + d) - \mathbf{m}_{\Delta}(x) = \mathbf{m}_{F_2(\Delta)}(x + d) - \mathbf{m}_{F_1(\Delta)}(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Let $s = \lambda m + \mu$ be the Euclidean division of s by m . Since $F_2(\Delta) = \text{AS}(a, (0, d), s)$, we know, from Lemma 3.5, that

$$\mathbf{m}_{F_2(\Delta)}(a + id) = \binom{\lambda + 1}{2} m + \left\lceil \frac{s - i}{m} \right\rceil (\mu - i)$$

for all $i \in [0, m-1]$. Moreover, since $F_1(\Delta) = \text{AS}(a, (d, d), s) = \text{AS}(a + (s-1)d, (0, -d), s)$, by Lemma 3.5 again, we have

$$\mathbf{m}_{F_1(\Delta)}(a + (s-1-j)d) = \binom{\lambda + 1}{2} m + \left\lceil \frac{s - j}{m} \right\rceil (\mu - j)$$

for all $j \in [0, m-1]$. This leads to the identity


$$\begin{aligned} \mathbf{m}_{\Delta}(a) - \mathbf{m}_{\Delta}(a - d) &= \mathbf{m}_{F_2(\Delta)}(a) - \mathbf{m}_{F_1(\Delta)}(a - d) \\ &= \mathbf{m}_{F_2(\Delta)}(a) - \mathbf{m}_{F_1(\Delta)}(a + (s-1-\mu)d) = \left\lceil \frac{s}{m} \right\rceil \mu. \end{aligned}$$

If Δ is balanced, then \mathbf{m}_{Δ} is constant on $\mathbb{Z}/m\mathbb{Z}$. Then, $\mathbf{m}_{\Delta}(a) = \mathbf{m}_{\Delta}(a - d)$ and $\left\lceil \frac{s}{m} \right\rceil \mu = 0$. Therefore $\mu = 0$ and $s \equiv 0 \pmod{m}$. Suppose now that $s = \lambda m$, with $\lambda \geq 1$. For every integer $i \in [1, m-1]$, we deduce from the previous results that

$$\begin{aligned} \mathbf{m}_{\Delta}(a + id) - \mathbf{m}_{\Delta}(a + (i-1)d) &= \mathbf{m}_{F_2(\Delta)}(a + id) - \mathbf{m}_{F_1(\Delta)}(a + (i-1)d) \\ &= \mathbf{m}_{F_2(\Delta)}(a + id) - \mathbf{m}_{F_1(\Delta)}(a + (s-1-(m-i))d) \\ &= \lambda(m-2i). \end{aligned}$$

The only possibility for having $\mathbf{m}_{\Delta}(a + id) = \mathbf{m}_{\Delta}(a + (i-1)d)$ is that $m = 2$ and $i = 1$. This concludes the proof. \square

Lemma 3.12. *Let s be a positive integer let $a, d \in \mathbb{Z}/4\mathbb{Z}$ such that d is invertible. Then, the arithmetic tetrahedron $\text{AS}(a, (0, d, -d), s)$ is not balanced in $\mathbb{Z}/4\mathbb{Z}$.*

FIGURE 8. $\text{AS}(a, (0, d, -d), s)$ in $\mathbb{Z}/4\mathbb{Z}$, with $d = \pm 1$

Proof. Let $\Delta = \text{AS}(a, (0, d, -d), s)$. If Δ is balanced in $\mathbb{Z}/4\mathbb{Z}$, then its projection $\pi_2(\Delta) = \text{AS}(\pi_2(a), (0, 1, 1), s)$ is balanced in $\mathbb{Z}/2\mathbb{Z}$. This implies, by Lemma 3.11, that s is even. Let $s = 4\lambda + \mu$ be the Euclidean division of s by 4, where $\mu \in \{0, 2\}$. Since $F_3(\Delta) = \text{AS}(a, (0, d), s)$ and $F_2(\Delta) = \text{AS}(a, (0, -d), s)$, it follows from Lemma 3.5 and Lemma 3.3 that

$$\begin{aligned} \mathbf{m}_\Delta(a) - \mathbf{m}_\Delta(a+2) &= \mathbf{m}_{F_3(\Delta)}(a) - \mathbf{m}_{F_2(\Delta)}(a+2) \\ &= \mathbf{m}_{F_3(\Delta)}(a) - \mathbf{m}_{F_2(\Delta)}(a-2d) \\ &= 4 \binom{\lambda+1}{2} + \left\lceil \frac{4\lambda+\mu}{4} \right\rceil \mu - 4 \binom{\lambda+1}{2} - \left\lceil \frac{4\lambda+\mu-2}{4} \right\rceil (\mu-2) \\ &= \left\lceil \frac{4\lambda+\mu}{4} \right\rceil \mu - \left\lceil \frac{4\lambda+\mu-2}{4} \right\rceil (\mu-2). \end{aligned}$$

For $\mu = 0$, we obtain that $\mathbf{m}_\Delta(a) - \mathbf{m}_\Delta(a+2) = 2\lambda \neq 0$. For $\mu = 2$, $\mathbf{m}_\Delta(a) - \mathbf{m}_\Delta(a+2) = 2(\lambda+1) \geq 2$. In all cases, we obtain that $\mathbf{m}_\Delta(a) \neq \mathbf{m}_\Delta(a+2)$. Therefore the tetrahedron Δ is not balanced in $\mathbb{Z}/4\mathbb{Z}$. \square

We are now ready to prove Theorem 3.8.

Proof of Theorem 3.8. Let $\Delta = \text{AS}(a, (d_1, d_2, d_3), s)$ be an arithmetic tetrahedron of size s , where all the common differences are

$$D := \{d_1, d_2, d_3, d_2 - d_1, d_3 - d_2, d_1 - d_3\}.$$

If there exists an element δ of D which is not invertible in $\mathbb{Z}/m\mathbb{Z}$ and not equal to zero, then we consider the projection of Δ into $\mathbb{Z}/\alpha\mathbb{Z}$, where $\alpha = \gcd(\delta, m) \geq 2$. The projected tetrahedron $\pi_\alpha(\Delta)$ is also arithmetic. In this tetrahedron, the corresponding common differences of the invertible common differences of Δ are invertible in $\mathbb{Z}/\alpha\mathbb{Z}$ and $\pi_\alpha(\delta) = 0$. If there exists a common difference δ' of $\pi_\alpha(D)$ which is not invertible in $\mathbb{Z}/\alpha\mathbb{Z}$ and not equal to zero, we project again $\pi_\alpha(\Delta)$ into $\mathbb{Z}/\beta\mathbb{Z}$, where $\beta = \gcd(\delta', \alpha) \geq 2$. We continue until the projected tetrahedron is such that all its common differences are either invertible, or equal to zero. In the sequel, suppose that $\pi_\alpha(\Delta)$ is like that. Obviously, if the projected tetrahedron $\pi_\alpha(\Delta)$ is not balanced in $\mathbb{Z}/\alpha\mathbb{Z}$, we know that Δ cannot be balanced in $\mathbb{Z}/m\mathbb{Z}$. We distinguish different cases.

Case 1. There exist three adjacent common differences, in $\pi_\alpha(\Delta)$, which are equal to zero. Then, all the elements of $\pi_\alpha(D)$ are equal to zero and $\pi_\alpha(\Delta)$ is the constant tetrahedron where all terms are equal to $\pi_\alpha(a)$. Thus, the tetrahedron $\pi_\alpha(\Delta)$ is not balanced in $\mathbb{Z}/\alpha\mathbb{Z}$.

Case 2. There exist two adjacent common differences, in $\pi_\alpha(\Delta)$, which are equal to zero. Without loss of generality, suppose that $\pi_\alpha(d_1) = \pi_\alpha(d_2) = 0$. Then, $\pi_\alpha(\Delta) =$

$\text{AS}(\pi_\alpha(a), (0, 0, \pi_\alpha(d_3)), s)$. If $\pi_\alpha(d_3) = 0$, this is Case 1. Otherwise, if $\pi_\alpha(d_3)$ is invertible, the tetrahedron $\pi_\alpha(\Delta)$ is not balanced in $\mathbb{Z}/\alpha\mathbb{Z}$, by Lemma 3.9.

Case 3. There exist two non adjacent common differences, in $\pi_\alpha(\Delta)$, which are equal to zero. Without loss of generality, suppose that $\pi_\alpha(d_1) = \pi_\alpha(d_3 - d_2) = 0$. Then, $\pi_\alpha(\Delta) = \text{AS}(\pi_\alpha(a), (0, \pi_\alpha(d_2), \pi_\alpha(d_2)), s)$. If $\pi_\alpha(d_2) = 0$, this is Case 1. Otherwise, if $\pi_\alpha(d_2)$ is invertible, we know from Lemma 3.11 that if $\pi_\alpha(\Delta)$ is balanced, then $\alpha = 2$ and s is even.

Case 4. There exists one common difference, in $\pi_\alpha(\Delta)$, which is equal to zero. Without loss of generality, suppose that $\pi_\alpha(d_1) = 0$. If there exists a second common difference which is equal to zero, this is Case 2 or Case 3, depending on the adjacency of these two common differences. If the five other common differences of $\pi_\alpha(\Delta)$ are invertible, we know that $\pi_\alpha(\Delta)$ is not balanced by Lemma 3.10.

Therefore, the only possibility for $\pi_\alpha(\Delta)$ being balanced in $\mathbb{Z}/\alpha\mathbb{Z}$ is that $\alpha = 2$ and s is even. Thus, when m is odd, we have proved the result i) of Theorem 3.8. When m is even, we deduce from the previous results that if Δ is balanced with a non invertible common difference δ , then the corresponding non adjacent common difference of δ is also non invertible and their projection in $\mathbb{Z}/2\mathbb{Z}$ are equal to zero. Moreover, since the constant tetrahedra in $\mathbb{Z}/2\mathbb{Z}$ are not balanced, it follows that the four other common differences of Δ must be invertible in $\mathbb{Z}/m\mathbb{Z}$. Suppose now that m and s are even and that the common differences of the balanced tetrahedron Δ are such that d_1 and $d_3 - d_2$ are non invertible and $d_2, d_3, d_2 - d_1, d_1 - d_3$ are invertible, where

$$\pi_2(\Delta) = \text{AS}(\pi_2(a), (0, 1, 1), s).$$

If $\gcd(d_1, m)$ or $\gcd(d_3 - d_2, m)$ are divisible by an odd number m' , then the projected tetrahedron $\pi_{m'}(\Delta)$ is a balanced arithmetic tetrahedron in $\mathbb{Z}/m'\mathbb{Z}$, with at least one common difference which is non invertible, that is impossible by result i) of Theorem 3.8. Thus $\gcd(d_1, m)$ and $\gcd(d_3 - d_2, m)$ are powers of two. Finally, from Lemma 3.12, we deduce that $\gcd(d_1, m) = \gcd(d_3 - d_2, m) = 2$. This completes the proof. \square

We continue by showing that there is no balanced arithmetic tetrahedron in $\mathbb{Z}/m\mathbb{Z}$ when m is divisible by 3.

Theorem 3.13. *Let m and s be two positive integer such that m is a multiple of 3. There is no balanced arithmetic tetrahedron of size s in $\mathbb{Z}/m\mathbb{Z}$.*

Proof. Let Δ be an arithmetic tetrahedron of size s in $\mathbb{Z}/m\mathbb{Z}$. We consider the projected tetrahedron $\pi_3(\Delta)$ in $\mathbb{Z}/3\mathbb{Z}$. We know from Theorem 3.8 that if $\pi_3(\Delta)$ is a balanced tetrahedron, then all the common differences of $\pi_3(\Delta)$ are invertible in $\mathbb{Z}/3\mathbb{Z}$. In $\mathbb{Z}/3\mathbb{Z}$, the invertible elements are 1 and 2. Thus, if d_1, d_2 and d_3 are invertible, there are at least two of them which are equal and thus their difference is a common difference equal to zero. Therefore $\pi_3(\Delta)$ cannot be balanced in $\mathbb{Z}/3\mathbb{Z}$. This completes the proof. \square

In the end of this subsection, we prove that the necessary conditions on the common differences of balanced arithmetic tetrahedra highlighted in Theorem 3.8 are also sufficient for certain sizes.

Theorem 3.14. *Let m be an odd number not divisible by 3 and let $a, d_1, d_2, d_3 \in \mathbb{Z}/m\mathbb{Z}$ such that $d_1, d_2, d_3, d_2 - d_1, d_3 - d_2$ and $d_1 - d_3$ are invertible. Then, the arithmetic tetrahedron $\text{AS}(a, (d_1, d_2, d_3), s)$ is balanced for all $s \equiv 0, -1, \text{ or } -2 \pmod{m}$.*

Proof. Theorem 3.2 for $n = 3$. \square

Theorem 3.15. *Let m be an even number not divisible by 3 and let $a, d_1, d_2, d_3 \in \mathbb{Z}/m\mathbb{Z}$ such that $\gcd(d_1, m) = \gcd(d_3 - d_2, m) = 2$ and $d_2, d_3, d_2 - d_1$ and $d_1 - d_3$ are invertible. Then, the arithmetic tetrahedron $\text{AS}(a, (d_1, d_2, d_3), s)$ is balanced for all $s \equiv 0$ or $-2 \pmod{m}$.*

Proof. Let $\Delta = \text{AS}(a, (d_1, d_2, d_3), s)$.

Case 1. Suppose that s is divisible by m .

Since $\gcd(d_1, m) = 2$, d_3 and $d_1 - d_3$ invertible and s divisible by m , it follows from Proposition 3.6 that the multiplicity function of $F_2(\Delta) = \text{AS}(a, (d_1, d_3), s)$ is entirely determined by

$$\mathbf{m}_{F_2(\Delta)}(a + 2i) = \frac{1}{m} \binom{s+1}{2} + \frac{s}{2m}, \quad \mathbf{m}_{F_2(\Delta)}(a + 2i + 1) = \frac{1}{m} \binom{s+1}{2} - \frac{s}{2m},$$

for all $i \in [0, \frac{m}{2} - 1]$. Similarly, since $\gcd(d_3 - d_2, m) = 2$, d_2 and d_3 invertible and s divisible by m , we have by Proposition 3.6 that the multiplicity function of $F_1(\Delta) = \text{AS}(a, (d_2, d_3), s) = \text{AS}(a + (s-1)d_2, d_3 - d_2, -d_2, s)$ is equal to

$$\mathbf{m}_{F_1(\Delta)}(a + 2i + 1) = \frac{1}{m} \binom{s+1}{2} + \frac{s}{2m}, \quad \mathbf{m}_{F_1(\Delta)}(a + 2i) = \frac{1}{m} \binom{s+1}{2} - \frac{s}{2m},$$

for all $i \in [0, \frac{m}{2} - 1]$. This leads to the identity

$$\mathbf{m}_{F_2(\Delta)}(x + 1) = \mathbf{m}_{F_1(\Delta)}(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Then, since $\gcd(d_1, m) = 2$ and d_2 is invertible, we obtain by Lemma 3.3 that

$$\begin{aligned} \mathbf{m}_{\Delta}(x + d_1) - \mathbf{m}_{\Delta}(x + d_2) &= \mathbf{m}_{F_1(\Delta)}(x + d_1) - \mathbf{m}_{F_2(\Delta)}(x + d_2) \\ &= \mathbf{m}_{F_1(\Delta)}(x + 2) - \mathbf{m}_{F_2(\Delta)}(x + 1) = 0, \end{aligned}$$

and thus

$$\mathbf{m}_{\Delta}(x + (d_1 - d_2)) = \mathbf{m}_{\Delta}(x)$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Since $d_1 - d_2$ is invertible, we conclude that \mathbf{m}_{Δ} is constant and Δ is balanced in $\mathbb{Z}/m\mathbb{Z}$.

Case 2. Now, suppose that $s \equiv -2 \pmod{m}$.

The tetrahedron Δ can be seen as the arithmetic tetrahedron Δ' of size $s + 2$ where the first two rows have been removed, i.e. $\Delta = \Delta' \setminus \{R_0, R_1\}$, where $\Delta' := \text{AS}(a - 2d_3, (d_1, d_2, d_3), s + 2)$, $R_0 := \text{AS}(a - 2d_3, (d_1, d_2), s + 2)$ and $R_1 := \text{AS}(a - d_3, (d_1, d_2), s + 1)$. Since $\gcd(d_1, m) = 2$, $d_2, d_2 - d_1, d_3$ invertible and $s + 2 \equiv 0$, $s + 1 \equiv -1 \pmod{m}$, it follows from Proposition 3.6 again that

$$\mathbf{m}_{R_0}(a + 2i) = \frac{1}{m} \binom{s+3}{2} + \frac{s+2}{2m}, \quad \mathbf{m}_{R_0}(a + 2i + 1) = \frac{1}{m} \binom{s+3}{2} - \frac{s+2}{2m},$$

$$\mathbf{m}_{R_1}(a + 2i + 1) = \frac{1}{m} \binom{s+2}{2} + \frac{s+2}{2m}, \quad \mathbf{m}_{R_1}(a + 2i) = \frac{1}{m} \binom{s+2}{2} - \frac{s+2}{2m}.$$

This leads to

$$\mathbf{m}_{R_0}(x) + \mathbf{m}_{R_1}(x) = \frac{1}{m} \left(\binom{s+3}{2} + \binom{s+2}{2} \right),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Therefore the multiset $R_0 \cup R_1$ is balanced in $\mathbb{Z}/m\mathbb{Z}$. Moreover, since $s + 2 \equiv 0 \pmod{m}$, we already know from Case 1 that Δ' is balanced. The multiset difference of balanced multisets is obviously balanced. This completes the proof. \square

Remark 3.16. When m is even not divisible by 3, if we suppose that $\gcd(d_1, m) = \gcd(d_3 - d_2, m) = 2$ and $d_2, d_3, d_2 - d_1$ and $d_1 - d_3$ are invertible, then the arithmetic tetrahedron $\text{AS}(a, (d_1, d_2, d_3), s)$ is not balanced for all $s \equiv -1 \pmod{m}$. Indeed, it can be seen as the multiset difference of the arithmetic tetrahedron $\text{AS}(a - d_3, (d_1, d_2, d_3), s + 1)$, which is balanced by Theorem 3.15, and the arithmetic triangle $\text{AS}(a - d_3, (d_1, d_2), s + 1)$, which is not balanced by Theorem 3.4.

4. BALANCED SIMPLICES GENERATED FROM ARITHMETIC ARRAYS

We are now ready to show that the orbits generated from arithmetic arrays by additive cellular automata are a source of balanced simplices.

4.1. The general case : in dimension $n \geq 2$. In this subsection, we prove Theorem 1.14, the main result of this paper, and, in corollary, the special case of the Pascal cellular automata is examined.

4.1.1. *For any ACA.* First, we recall Theorem 1.14.

Theorem 4.1. *Let $n \geq 2$ and m be two positive integers such that $\gcd(m, n!) = 1$. Suppose that σ is invertible modulo m . Let $a \in \mathbb{Z}/m\mathbb{Z}$, $d = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$ and $\varepsilon \in \{-1, 1\}^n$ such that d_i , for all $1 \leq i \leq n$, and $\varepsilon_j d_j - \varepsilon_i d_i$, for all $1 \leq i < j \leq n$, are invertible, where $d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k$. Then, in the orbit $\mathcal{O}(\text{AA}(a, d))$, every n -simplex with orientation ε and of size s is balanced, for all $s \equiv -t \pmod{\text{ord}_m(\sigma^m)m}$, where $t \in [0, n - 1]$.*

The proof of this theorem is based on the following lemma.

Lemma 4.2. *Let $n \geq 2$, m_1 and m_2 be three positive integers such that $\gcd(m_2, n!) = 1$ and let $m := m_1 m_2$. Let $a \in \mathbb{Z}/m\mathbb{Z}$ and let $d = (d_1, \dots, d_n) \in (\mathbb{Z}/m\mathbb{Z})^n$ such that $\pi_{m_2}(d_i)$, for all $1 \leq i \leq n$, and $\pi_{m_2}(d_j) - \pi_{m_2}(d_i)$, for all $1 \leq i < j \leq n$, are invertible in $\mathbb{Z}/m_2\mathbb{Z}$. Then, for all positive integers $s \equiv -t \pmod{m_2}$, where $t \in [0, n - 1]$, the multiplicity function of the arithmetic simplex $\Delta = \text{AS}(a, m_1 d, s)$ of $\mathbb{Z}/m\mathbb{Z}$ verifies*

$$\mathbf{m}_\Delta(x + m_1) = \mathbf{m}_\Delta(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$.

Proof. First, it is clear that Δ is uniquely constituted by elements of the form $a + km_1$, where $k \in [0, m_2 - 1]$. Therefore, the identity is obviously true for all elements x not in $\{a + km_1 \mid k \in [0, m_2 - 1]\}$. Moreover, since $\pi_{m_2}(d_i)$, for all $1 \leq i \leq n$, and $\pi_{m_2}(d_j) - \pi_{m_2}(d_i)$, for all $1 \leq i < j \leq n$, are invertible in $\mathbb{Z}/m_2\mathbb{Z}$ and since $s \equiv -t \pmod{m_2}$, where $t \in [0, n - 1]$, we know from Theorem 3.2 that the arithmetic simplex $\text{AS}(0, \pi_2(d), s)$ is balanced in $\mathbb{Z}/m_2\mathbb{Z}$. Finally, since Δ can be seen as the image of the balanced arithmetic simplex $\Delta' = \text{AS}(0, \pi_2(d), s)$ by the function

$$\begin{aligned} f : \mathbb{Z}/m_2\mathbb{Z} &\longrightarrow \mathbb{Z}/m\mathbb{Z} \\ \pi_{m_2}(x) &\longmapsto a + m_1 x \end{aligned}$$

it follows that

$$\mathbf{m}_\Delta(a + km_1) = \mathbf{m}_{f(\Delta')}(f(k)) = \mathbf{m}_{\Delta'}(k) = \mathbf{m}_{\Delta'}(0) = \mathbf{m}_{f(\Delta')}(f(0)) = \mathbf{m}_\Delta(a),$$

for all $k \in [0, m_2 - 1]$. This completes the proof. \square

We are now ready to prove Theorem 4.1 (Theorem 1.14), the main result of this paper.

Proof of Theorem 4.1. Let $\Delta(j, \varepsilon, s)$ be an n -simplex of size $s = \lambda \text{lcm}(\text{ord}_m(\sigma), m) - t$, where $t \in [0, n - 1]$, appearing in the orbit $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^{n-1} \times \mathbb{N}}$. We proceed by induction on m . For $m = 1$, the result is obvious. Suppose now that the result is true for all finite cyclic groups of order strictly lesser than m . Let

$$m_1 := \gcd(\text{ord}_m(\sigma), m) \quad \text{and} \quad m_2 := \frac{m}{m_1}.$$

Then, $s = \lambda \text{ord}_m(\sigma) m_2 - t$.

First, we prove that $\mathbf{m}_\Delta(x + m_1) = \mathbf{m}_\Delta(x)$ for all $x \in \mathbb{Z}/m\mathbb{Z}$. By Proposition 2.11 for $\alpha = \text{ord}_m(\sigma)$, we know that $\Delta(j, \varepsilon, s)$ can be decomposed into $\text{ord}_m(\sigma)^n$ subsimplices SS_k ,

$$\Delta(j, \varepsilon, s) = \bigcup_{k \in [0, \text{ord}_m(\sigma) - 1]^n} \text{SS}_k$$

that are, for all $k \in [0, \text{ord}_m(\sigma) - 1]^n$, the arithmetic simplices

$$\text{SS}_k = \text{AS} \left(a_{j+\varepsilon \cdot k}, \text{ord}_m(\sigma) \sigma^{j_n + \varepsilon_n k_n} \varepsilon \cdot \tilde{d}, \lambda m_2 - \left\lfloor \frac{\sum_{u=1}^n k_u + t}{\text{ord}_m(\sigma)} \right\rfloor \right),$$

where $\tilde{d} = (d_1, \dots, d_{n-1}, d_n)$ and $d_n = \sigma^{-1} \sum_{u=1}^{n-1} \sigma_u d_u$. Since

$$\gcd\left(\frac{\text{ord}_m(\sigma)}{m_1}, m_2\right) = \frac{\gcd(\text{ord}_m(\sigma), m)}{\gcd(\text{ord}_m(\sigma), m)} = 1,$$

it follows, from the hypothesis of the theorem, that the elements

$$\pi_{m_2} \left(\frac{\text{ord}_m(\sigma)}{m_1} \sigma^{j_n + \varepsilon_n k_n} \varepsilon_u d_u \right),$$

for all $1 \leq u \leq n$, and

$$\pi_{m_2} \left(\frac{\text{ord}_m(\sigma)}{m_1} \sigma^{j_n + \varepsilon_n k_n} (\varepsilon_u d_u - \varepsilon_v d_v) \right),$$

for all $1 \leq u < v \leq n$, are invertible in $\mathbb{Z}/m_2\mathbb{Z}$. Moreover, since the size of SS_k , $\lambda m_2 - \left\lfloor \frac{\sum_{u=1}^n k_u + t}{\text{ord}_m(\sigma)} \right\rfloor$, is in $[\lambda m_2 - (n - 1), \lambda m_2]$ and thus is congruent to a certain integer $-y$ modulo m_2 with $y \in [0, n - 1]$, we deduce from Lemma 4.2 that

$$\mathbf{m}_{\text{SS}_k}(x + m_1) = \mathbf{m}_{\text{SS}_k}(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$, and this, for all $k \in [0, \text{ord}_m(\sigma) - 1]^n$. Then, the multiplicity function of Δ verifies

$$\mathbf{m}_\Delta(x + m_1) = \sum_{k \in [0, \text{ord}_m(\sigma) - 1]^n} \mathbf{m}_{\text{SS}_k}(x + m_1) = \sum_{k \in [0, \text{ord}_m(\sigma) - 1]^n} \mathbf{m}_{\text{SS}_k}(x) = \mathbf{m}_\Delta(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$.

Now, we prove that the projected simplex $\pi_{m_1}(\Delta)$ is balanced. We begin by showing that the integer $\text{ord}_m(\sigma^m)m$ is divisible by $\text{ord}_{m_1}(\sigma^{m_1})m_1$. Indeed,

$$(\sigma^{m_1})^{m_2 \text{ord}_m(\sigma^m)} = (\sigma^m)^{\text{ord}_m(\sigma^m)} \equiv 1 \pmod{m}$$

and, by divisibility of m by m_1 , this equivalence is also true modulo m_1 . It follows that $m_2 \text{ord}_m(\sigma^m)$ is divisible by $\text{ord}_{m_1}(\sigma^{m_1})$, implying that $\text{ord}_m(\sigma^m)m$ is divisible by $\text{ord}_{m_1}(\sigma^{m_1})m_1$. Then, $s \equiv -t \pmod{\text{ord}_{m_1}(\sigma^{m_1})m_1}$ and since the elements $\pi_{m_1}(d_u)$, for all $1 \leq u \leq n$, and $\pi_{m_1}(\varepsilon_v d_v - \varepsilon_u d_u)$, for all $1 \leq u < v \leq n$, are clearly invertible in $\mathbb{Z}/m_1\mathbb{Z}$, we deduce from the induction hypothesis that $\pi_{m_1}(\Delta)$ is balanced in $\mathbb{Z}/m_1\mathbb{Z}$.

Finally, since $\pi_{m_1}(\Delta)$ is balanced and $\mathbf{m}_\Delta(x + m_1) = \mathbf{m}_\Delta(x)$ for all $x \in \mathbb{Z}/m\mathbb{Z}$, we deduce from Theorem 2.5 that the simplex Δ is balanced. This concludes the proof. \square

Remark 4.3. If $\sigma \equiv 1 \pmod{m}$, then the n -simplex $\Delta(j, \varepsilon, s)$ appearing in $\mathcal{O}(\text{AA}(a, d))$ is an arithmetic simplex and Theorem 4.1 is simply Theorem 3.2 on balanced arithmetic simplices.

4.1.2. *For the Pascal cellular automata.* Here, we investigate the consequences of Theorem 4.1 on the existence of balanced n -simplices, in the case where the ACA considered is PCA_{n-1} .

Corollary 4.4. *Let $n \geq 2$ be a positive integers. For every positive integer m such that $\gcd(m, (3(n-1))!) = 1$, there exist infinitely many balanced n -simplices of $\mathbb{Z}/m\mathbb{Z}$ generated by PCA_{n-1} , for all possible orientations $\varepsilon \in \{-1, 1\}^n$. In the special case of the two orientations $\varepsilon = +\cdots+-$ or $\varepsilon = -\cdots-+$, the existence of an infinite number of such balanced simplices is verified for every $\mathbb{Z}/m\mathbb{Z}$ such that $\gcd(m, n!) = 1$, if n is even, and for every $\mathbb{Z}/m\mathbb{Z}$ such that $\gcd(m, (\frac{3n+1}{2})!) = 1$, if n is odd.*

Proof. For the Pascal automaton of dimension $n-1$, we have $\sigma = n$ and $\sigma_k = -1$ for all $k \in [1, n-1]$. Let m be a positive integer such that $\gcd(m, n!) = 1$ and let $A = \text{AA}(a, d)$ be the arithmetic array of $\mathbb{Z}/m\mathbb{Z}$ with common difference $d = (d_1, \dots, d_{n-1})$ defined by $d_k := k \in \mathbb{Z}/m\mathbb{Z}$ for all $k \in [1, n-1]$. Then,

$$d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k = -n^{-1} \sum_{k=1}^{n-1} k = -2^{-1}(n-1).$$

Since $\gcd(m, n!) = 1$, the common differences d_1, d_2, \dots, d_{n-1} and d_n are invertible in $\mathbb{Z}/m\mathbb{Z}$. For all integers u and v such that $1 \leq u < v \leq n-1$, we have

$$(1) \quad 3 \leq u+v \leq 2n-3 \quad \text{and} \quad 1 \leq v-u \leq n-2.$$

For all integers u such that $1 \leq u \leq n-1$, we have

$$(2) \quad n+1 \leq 2u+(n-1) \leq 3(n-1) \quad \text{and} \quad -(n-3) \leq 2u-(n-1) \leq n-1.$$

If n is even, then $2u-(n-1)$ cannot vanish and we deduce that if $\gcd(m, (3(n-1))!) = 1$ then, for any orientation $\varepsilon \in \{-1, 1\}^n$, all the elements in

$$(3) \quad \{\varepsilon_v d_v - \varepsilon_u d_u \mid 1 \leq u < v \leq n\}$$

are invertible modulo m . Therefore, by Theorem 4.1, any simplices of orientation ε and of size $s \equiv -t \pmod{\text{lcm}(\text{ord}_m(n), m)}$, with $t \in [0, n-1]$, appearing in the orbit $\mathcal{O}(A)$ are balanced in $\mathbb{Z}/m\mathbb{Z}$. For the specific orientations $\varepsilon = +\cdots+-$ and $\varepsilon = -\cdots-+$, we deduce from the second inequalities of (1) and (2) that this result is also true in the more general case where $\gcd(m, n!) = 1$. Now, suppose that n is odd and let $\varepsilon \in \{-1, 1\}^n$ be an orientation. If there exist $l \in [1, n-1]$ such that $\varepsilon_l = \varepsilon_n$, then we consider the arithmetic array $A = \text{AA}(a, d)$ of $\mathbb{Z}/m\mathbb{Z}$ with common differences $d = (d_1, \dots, d_{n-1})$ defined by $d_l := \frac{n-1}{2}$, $d_{\frac{n-1}{2}} := l$ and $d_k := k$ for all $k \in [1, n-1] \setminus \{l, \frac{n-1}{2}\}$. Then, $d_n = -2^{-1}(n-1)$ as before and

$$\varepsilon_l d_l - \varepsilon_n d_n = \varepsilon_l(n-1).$$

If $\gcd(m, (3(n-1))!) = 1$, we deduce that all the common differences d_k for $k \in [1, n]$ and, from inequalities (1) and (2), all the elements in the set of (3) are invertible modulo m . Therefore, by Theorem 4.1, any simplices of this orientation ε , where $\varepsilon_l = \varepsilon_n$, and of size $s \equiv -t \pmod{\text{lcm}(\text{ord}_m(n), m)}$, with $t \in [0, n-1]$, appearing in the orbit $\mathcal{O}(A)$ are balanced in $\mathbb{Z}/m\mathbb{Z}$. Finally, suppose that n is odd and ε is such that $\varepsilon_k = -\varepsilon_n$ for all $k \in [1, n-1]$, i.e., $\varepsilon = +\cdots+-$ or $\varepsilon = -\cdots-+$. In this case, we consider the arithmetic array $A = \text{AA}(a, d)$ of $\mathbb{Z}/m\mathbb{Z}$ with common differences $d = (d_1, \dots, d_{n-1})$ defined by

$d_{\frac{n+1}{2}} := \frac{3n+1}{2}$ and $d_k := k$ for all $k \in [1, n-1] \setminus \{\frac{n+1}{2}\}$. Then, $d_n = -2^{-1}(n+1)$. For all integers u and v such that $1 \leq u < v \leq \frac{3n+1}{2}$, we have

$$(4) \quad 1 \leq v - u \leq \frac{3n-1}{2} \quad \text{and} \quad -\frac{n-1}{2} \leq u - \frac{n+1}{2} \leq n.$$

If $\mathbb{Z}/m\mathbb{Z}$ is such that $\gcd(m, (\frac{3n+1}{2})!) = 1$, we know by definition that all the common differences d_k are invertible modulo m for all $k \in [1, n]$. Moreover, since $d_u + d_n$ cannot vanish by definition, and from (4), we deduce that all the elements in the set of (3) are invertible modulo m . By Theorem 4.1 again, any simplices with these orientations, $\varepsilon = +\cdots+-$ or $\varepsilon = -\cdots-+$, and of size $s \equiv -t \pmod{\text{lcm}(\text{ord}_m(n), m)}$, with $t \in [0, n-1]$, appearing in the orbit $\mathcal{O}(A)$ are balanced in $\mathbb{Z}/m\mathbb{Z}$. This concludes the proof. \square

4.2. In dimension 3. In this subsection, we show that, in dimension 3, a similar result than Theorem 4.1 can be obtained for certain even values of m by using Theorem 3.15. In corollary, the special case of the Pascal cellular automaton PCA_2 is studied.

4.2.1. *For any ACA.*

Theorem 4.5. *Let m be an even number not divisible by 3 such that $\sigma \in \mathbb{Z}/m\mathbb{Z}$ is invertible and $\sigma \equiv 1 \pmod{2^{v_2(m)}}$, where $v_2(m)$ is the highest exponent u such that 2^u divides m . Let $a \in \mathbb{Z}/m\mathbb{Z}$, $d = (d_1, d_2) \in (\mathbb{Z}/m\mathbb{Z})^2$ and $\varepsilon \in \{-1, 1\}^3$ such that $\varepsilon_2 d_2, \varepsilon_3 d_3, \varepsilon_2 d_2 - \varepsilon_1 d_1, \varepsilon_1 d_1 - \varepsilon_3 d_3$ are invertible in $\mathbb{Z}/m\mathbb{Z}$ and $\gcd(\varepsilon_1 d_1, m) = \gcd(\varepsilon_3 d_3 - \varepsilon_2 d_2, m) = 2$, where $d_3 := \sigma^{-1}(\sigma_1 d_1 + \sigma_2 d_2)$. Then, in the orbit $\mathcal{O}(\text{AA}(a, d))$, every tetrahedron with orientation ε and of size s is balanced, for all $s \equiv 0$ or $-2 \pmod{\text{lcm}(\text{ord}_m(\sigma), m)}$.*

Proof. Let $\Delta = \Delta(j, \varepsilon, s)$ be a tetrahedron of size $\lambda \text{lcm}(\text{ord}_m(\sigma), m) - t$, where $t \in \{0, 2\}$, appearing in the orbit $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^2 \times \mathbb{N}}$. We proceed by induction on m .

For $m = 2^{v_2(m)}$, since $\sigma \equiv 1 \pmod{m}$, it follows that Δ is an arithmetic tetrahedron of size $s \equiv 0$ or $-2 \pmod{m}$ and of common differences $(\varepsilon_1 d_1, \varepsilon_2 d_2, \varepsilon_3 d_3)$ such that $\varepsilon_2 d_2, \varepsilon_2 d_2 - \varepsilon_1 d_1, \varepsilon_3 d_3$ and $\varepsilon_1 d_1 - \varepsilon_3 d_3$ are invertible and $\gcd(\varepsilon_1 d_1, m) = \gcd(\varepsilon_3 d_3 - \varepsilon_2 d_2) = 2$. By Theorem 3.15, the tetrahedron Δ is balanced in this case.

Suppose now that $m > 2^{v_2(m)}$ and let $m' := m/2^{v_2(m)}$, the odd part of m . Let

$$m_1 := 2^{v_2(m)} \gcd(\text{ord}_m(\sigma), m') \quad \text{and} \quad m_2 := \frac{m}{m_1} = \frac{m'}{\gcd(\text{ord}_m(\sigma), m')}.$$

First, we prove that $\mathbf{m}_\Delta(x + m_1) = \mathbf{m}_\Delta(x)$, for all $x \in \mathbb{Z}/m\mathbb{Z}$. Since

$$m_1 = 2^{v_2(m)} \gcd(\text{ord}_m(\sigma), m') = \gcd(2^{v_2(m)} \text{ord}_m(\sigma), m),$$

it follows that m_1 is divisible by $\gcd(\text{ord}_m(\sigma), m)$. By Proposition 2.11 with

$$\alpha = \frac{m_1}{\gcd(\text{ord}_m(\sigma), m)} \text{ord}_m(\sigma),$$

since $s = \alpha m_2 - t$, we know that Δ can be decomposed into α^3 subtetrahedra that are arithmetic. More precisely, we have

$$\Delta = \bigcup_{k \in [0, \alpha-1]^3} \text{SS}_k,$$

where

$$\text{SS}_k = \text{AS} \left(a_{j+\varepsilon \cdot k}, m_1 \frac{\text{ord}_m(\sigma)}{\gcd(\text{ord}_m(\sigma), m)} \sigma^{j_3 + \varepsilon_3 k_3} (\varepsilon_1 d_1, \varepsilon_2 d_2, \varepsilon_3 d_3), m_2 - \left\lfloor \frac{k_1 + k_2 + k_3}{\alpha} \right\rfloor \right),$$

for all $k \in [0, \alpha - 1]^3$. Since m_2 is an odd factor of m and $\gcd(\varepsilon_2 d_2, m) = \gcd(\varepsilon_3 d_3 - \varepsilon_2 d_2, m) = 2$, we deduce that $\pi_{m_2}(\varepsilon_2 d_2)$ and $\pi_{m_2}(\varepsilon_3 d_3 - \varepsilon_2 d_2)$ are invertible in $\mathbb{Z}/m_2\mathbb{Z}$. Since σ , $\varepsilon_2 d_2$, $\varepsilon_3 d_3$, $\varepsilon_2 d_2 - \varepsilon_1 d_1$ and $\varepsilon_1 d_1 - \varepsilon_3 d_3$ are invertible in $\mathbb{Z}/m\mathbb{Z}$ and since

$$\gcd\left(\frac{\text{ord}_m(\sigma)}{\gcd(\text{ord}_m(\sigma), m)}, m_2\right) \mid \gcd\left(\frac{\text{ord}_m(\sigma)}{\gcd(\text{ord}_m(\sigma), m')}, \frac{m'}{\gcd(\text{ord}_m(\sigma), m')}\right) = 1,$$

we obtain that the elements

$$\pi_{m_2}\left(\frac{\text{ord}_m(\sigma)}{\gcd(\text{ord}_m(\sigma), m)}\sigma^{j_3+\varepsilon_3 k_3}\varepsilon_u d_u\right),$$

for all $1 \leq u \leq 3$, and

$$\pi_{m_2}\left(\frac{\text{ord}_m(\sigma)}{\gcd(\text{ord}_m(\sigma), m)}\sigma^{j_3+\varepsilon_3 k_3}(\varepsilon_v d_v - \varepsilon_u d_u)\right),$$

for all $1 \leq u < v \leq 3$, are invertible in $\mathbb{Z}/m_2\mathbb{Z}$. Moreover, since the size of SS_k is $m_2 - \lfloor \frac{k_1+k_2+k_3}{\alpha} \rfloor$, which is congruent to 0, -1 , or -2 modulo m_2 , we deduce from Lemma 4.2 that the multiplicity function of SS_k verifies

$$\mathbf{m}_{\text{SS}_k}(x + m_1) = \mathbf{m}_{\text{SS}_k}(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$ and for all $k \in [0, \alpha - 1]^3$. Therefore, we have

$$\mathbf{m}_\Delta(x + m_1) = \sum_{k \in [0, \alpha - 1]^3} \mathbf{m}_{\text{SS}_k}(x + m_1) = \sum_{k \in [0, \alpha - 1]^3} \mathbf{m}_{\text{SS}_k}(x) = m_\Delta(x),$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Since $\sigma \equiv 1 \pmod{2^{v_2(m)}}$, it follows that $\text{ord}_m(\sigma) = \text{ord}_{m'}(\sigma)$ and thus $m_1 = 2^{v_2(m)} \gcd(\text{ord}_m(\sigma), m') = 2^{v_2(m)} \gcd(\text{ord}_{m'}(\sigma), m')$ is a proper divisor of m . Thus, from the induction hypothesis, we know that the projection of Δ into $\mathbb{Z}/m_1\mathbb{Z}$ is balanced. Finally, since $\mathbf{m}_\Delta(x + m_1) = \mathbf{m}_\Delta(x)$ for all $x \in \mathbb{Z}/m\mathbb{Z}$ and $\pi_{m_1}(\Delta)$ balanced in $\mathbb{Z}/m_1\mathbb{Z}$, we conclude that the tetrahedron Δ is balanced in $\mathbb{Z}/m\mathbb{Z}$ by Theorem 2.5. This completes the proof. \square

4.2.2. For the Pascal cellular automaton. Here, we investigate the consequences of Theorem 4.5 on the existence of balanced tetrahedra, in the case where the ACA considered is PCA_2 .

Corollary 4.6. *For even numbers m not divisible by 3 and such that $v_2(m) = 1$, there exist infinitely many balanced tetrahedra of $\mathbb{Z}/m\mathbb{Z}$ generated by PCA_2 , for all orientations $\varepsilon = + + +$, $+ - +$, $+ - -$, $- + +$, $- + -$ and $- - -$. In the special case of the two orientations $\varepsilon = + + -$ or $\varepsilon = - - +$, the existence of an infinite number of such balanced tetrahedra is verified for every $\mathbb{Z}/m\mathbb{Z}$ of even order m such that $v_2(m) = 1$ and $\gcd(m, 3.5) = 1$.*

Proof. For the Pascal automaton of dimension 2, we have $\sigma = 3$ and $\sigma_1 = \sigma_2 = -1$. If there exists $i \in \{1, 2\}$ such that $\varepsilon_i = \varepsilon_3$, then we consider the orbit associated with the arithmetic array $\text{AA}(a, (d_1, d_2))$ of $\mathbb{Z}/m\mathbb{Z}$, where $d_i := 1$ and $d_j := 2$ with $\{i, j\} = \{1, 2\}$. Then, $d_3 = -3^{-1} \cdot 3 = -1$, $d_i + d_j = 3$, $d_j - d_i = 1$, $d_j + d_3 = 1$, $d_j - d_3 = 3$ and $d_i - d_3 = 2$. It follows that, if m is an even number not divisible by 3 and such that $v_2(m) = 1$, we have

$$\gcd(\varepsilon_i d_i, m) = \gcd(\varepsilon_3 d_3, m) = \gcd(\varepsilon_j d_j - \varepsilon_i d_i, m) = \gcd(\varepsilon_j d_j - \varepsilon_3 d_3, m) = 1$$

and

$$\gcd(\varepsilon_j d_j, m) = \gcd(\varepsilon_i d_i - \varepsilon_3 d_3, m) = 2.$$

Therefore, by Theorem 4.5, any tetrahedron of this orientation ε , where $\varepsilon_i = \varepsilon_3$, and of size $s \equiv 0$ or $-2 \pmod{\text{lcm}(\text{ord}_m(3), m)}$ appearing in the orbit $\mathcal{O}(A)$ is balanced in $\mathbb{Z}/m\mathbb{Z}$.

0	1	1	3	6	0	4	0	1	0	2	3	0	5	0
6	0	4	5	3	0	3	5	0	4	2				
6	3	0					0	4						
4	2						6							
1														

FIGURE 9. An $(1, 2)$ -antisymmetric tetrahedron in $\mathbb{Z}/7\mathbb{Z}$

Suppose now that $\varepsilon = ++-$ or $--+$ and consider the orbit associated with the arithmetic array $AA(a, (d_1, d_2))$ of $\mathbb{Z}/m\mathbb{Z}$, where $d_1 := 4$ and $d_2 := 5$. Then, $d_3 = -3^{-1} \cdot 9 = -3$, $d_2 - d_1 = 1$, $d_1 + d_3 = 1$ and $d_2 + d_3 = 2$. It follows that, if m is an even number such that $v_2(m) = 1$ and $\gcd(m, 3 \cdot 5) = 1$, we have

$$\gcd(\varepsilon_2 d_2, m) = \gcd(\varepsilon_3 d_3, m) = \gcd(\varepsilon_2 d_2 - \varepsilon_1 d_1, m) = \gcd(\varepsilon_1 d_1 - \varepsilon_3 d_3, m) = 1$$

and

$$\gcd(\varepsilon_1 d_1, m) = \gcd(\varepsilon_2 d_2 - \varepsilon_3 d_3, m) = 2.$$

Therefore, by Theorem 4.5, any tetrahedron of the orientation $\varepsilon = ++-$ or $--+$ and of size $s \equiv 0$ or $-2 \pmod{\text{lcm}(\text{ord}_m(3), m)}$ appearing in the orbit $\mathcal{O}(A)$ is balanced in $\mathbb{Z}/m\mathbb{Z}$. This concludes the proof. \square

5. THE ANTISYMMETRIC CASE

We begin this section by defining the antisymmetric sequences and the antisymmetric simplices.

Definition 5.1 (Antisymmetric sequences). A finite sequence $S = (a_1, \dots, a_s)$ of length $s \geq 1$ in $\mathbb{Z}/m\mathbb{Z}$ is said to be *antisymmetric* if $a_i + a_{s-i+1} = 0$ for all $i \in [1, s]$.

For instance, the sequence $S = (2, 2, 1, 0, 4, 3, 3)$ is antisymmetric in $\mathbb{Z}/5\mathbb{Z}$.

Definition 5.2 (Antisymmetric simplices). Let $A = (a_i)_{i \in \mathbb{Z}^n}$ be an infinite array of elements in $\mathbb{Z}/m\mathbb{Z}$ and let $\Delta(j, \varepsilon, s)$ be the n -simplex of size s , with orientation $\varepsilon \in \{-1, 1\}^n$ and whose principal vertex is a_j in A , that is,

$$\Delta(j, \varepsilon, s) = \{a_{j+\varepsilon \cdot k} \mid k \in \mathbb{N}^n \text{ such that } k_1 + \dots + k_n \leq s - 1\}.$$

Let u and v be two integers such that $0 \leq u < v \leq n$. The simplex $\Delta(j, \varepsilon, s)$ is said to be (u, v) -*antisymmetric* if all its subsequences in the same direction of the edge between the vertices V_u and V_v are antisymmetric. More precisely, $\Delta(j, \varepsilon, s)$ is $(0, v)$ -antisymmetric if we have

$$a_{j+\varepsilon \cdot k} + a_{j+\varepsilon \cdot s(k)} = 0, \text{ where } s(k) = \left(k_1, \dots, k_{v-1}, s - 1 - \sum_{l=1}^n k_l, k_{v+1}, \dots, k_n \right),$$

for all $k \in \mathbb{N}^n$ such that $k_1 + \dots + k_n \leq s - 1$ and, for $u \geq 1$, $\Delta(j, \varepsilon, s)$ is (u, v) -antisymmetric if we have

$$a_{j+\varepsilon \cdot k} + a_{j+\varepsilon \cdot t(k)} = 0, \text{ where } (t(k))_l = k_{\tau(l)} \text{ for all } l \in [1, n],$$

where τ is the transposition (u, v) , for all $k \in \mathbb{N}^n$ such that $k_1 + \dots + k_n \leq s - 1$.

For instance, the tetrahedron depicted in Figure 9 is $(1, 2)$ -antisymmetric. Moreover, each row of this tetrahedron is an $(1, 2)$ -antisymmetric triangle.

5.1. Antisymmetric simplices. Let m and n be two positive integers such that $n \geq 2$ and $\gcd(m, n!) = 1$. In the sequel of this section, we consider n -simplices $\Delta(j, \varepsilon, s)$ appearing in the orbit of the arithmetic array $\text{AA}(a, d)$, where $a \in \mathbb{Z}/m\mathbb{Z}$ and $d = (d_1, d_2, \dots, d_{n-1}) \in (\mathbb{Z}/m\mathbb{Z})^{n-1}$, generated by an ACA of weight array $W = (w_l)_{l \in [-r, r]^{n-1}}$. The elements of this orbit are denoted by $\mathcal{O}(\text{AA}(a, d)) = (a_i)_{i \in \mathbb{Z}^{n-1} \times \mathbb{N}}$. As already defined before,

$$\sigma := \sum_{l \in [-r, r]^{n-1}} w_l \quad \text{and} \quad \sigma_k = \sum_{l \in [-r, r]^{n-1}} l_k w_l, \quad \text{for all } k \in [1, n-1].$$

Moreover, suppose that σ is invertible modulo m and let

$$d_n := \sigma^{-1} \sum_{k=1}^{n-1} \sigma_k d_k.$$

In this subsection, necessary conditions on simplices for being antisymmetric are determined.

Proposition 5.3. *Let u and v be two integers such that $1 \leq u < v \leq n$. If $\Delta(j, \varepsilon, s)$ is (u, v) -antisymmetric, then $d_w = 0$ for all $w \in [1, n] \setminus \{u, v\}$ and $\varepsilon_u d_u + \varepsilon_v d_v = 0$.*

Proof. Let $k \in \mathbb{N}^n$ such that $k_1 + \dots + k_n \leq s - 1$. If $k_u = k_v$, then $\tau(k) = k$ and

$$2a_{j+\varepsilon \cdot k} = a_{j+\varepsilon \cdot k} + a_{j+\varepsilon \cdot \tau(k)} = 0,$$

by definition of the (u, v) -antisymmetry. It follows that $a_{j+\varepsilon \cdot k} = 0$ in this case. Therefore, if $k_u = k_v$, we have

$$\sigma^{j_n + \varepsilon_n k_n} \left(a + \sum_{l=1}^n (j_l + \varepsilon_l k_l) d_l \right) = 0 \iff a + \sum_{l=1}^n (j_l + \varepsilon_l k_l) d_l = 0.$$

For $k = 0$, we obtain that

$$a + \sum_{l=1}^n j_l d_l = 0.$$

Now, we consider the canonical basis (e_1, \dots, e_n) of the vector space \mathbb{Z}^n . For all $w \in [1, n] \setminus \{u, v\}$, since $k_u = k_v = 0$ in $k = e_w$, we have

$$a_{j+\varepsilon \cdot e_w} = 0 \iff a + \sum_{l=1}^n j_l d_l + \varepsilon_w d_w = 0 \iff \varepsilon_w d_w = 0 \iff d_w = 0.$$

Finally, for $k = e_u + e_v$, since $k_u = k_v = 1$, we obtain that

$$a_{j+\varepsilon \cdot k} = 0 \iff a + \sum_{l=1}^n j_l d_l + \varepsilon_u d_u + \varepsilon_v d_v = 0 \iff \varepsilon_u d_u + \varepsilon_v d_v = 0.$$

This completes the proof. □

Proposition 5.4. *Let v be an integer in $[1, n]$. If $\Delta(j, \varepsilon, s)$ is $(0, v)$ -antisymmetric, then $2\varepsilon_w d_w = \varepsilon_v d_v$ for all $w \in [1, n] \setminus \{v\}$.*

Proof. Let $k \in \mathbb{N}^n$ such that $k_1 + \dots + k_n \leq s - 1$. If $k_v = s - 1 - \sum_{l=1}^n k_l$, then $s(k) = k$ and

$$2a_{j+\varepsilon \cdot k} = a_{j+\varepsilon \cdot k} + a_{j+\varepsilon \cdot s(k)} = 0,$$

by definition of the $(0, v)$ -antisymmetry. It follows that $a_{j+\varepsilon \cdot k} = 0$ in this case. Therefore, if $k_v = s - 1 - \sum_{l=1}^n k_l$, we have

$$\sigma^{j_n + \varepsilon_n k_n} \left(a + \sum_{l=1}^n (j_l + \varepsilon_l k_l) d_l \right) = 0 \iff a + \sum_{l=1}^n (j_l + \varepsilon_l k_l) d_l = 0.$$

Let $w \in [1, n] \setminus \{v\}$. Since $k_v + \sum_{l=1}^n k_l = s - 1$ for $k = (s - 1)e_w$, we have

$$a_{j+\varepsilon \cdot (s-1)e_w} = 0 \iff a + \sum_{l=1}^n j_l d_l + \varepsilon_w (s-1)d_w = 0.$$

Moreover, for $k = e_v + (s - 3)e_w$, since $\sum_{l=1}^n k_l + k_v = s - 1$, we obtain that

$$a_{j+\varepsilon \cdot k} = 0 \iff a + \sum_{l=1}^n j_l d_l + \varepsilon_v d_v + \varepsilon_w (s-3)d_w = 0.$$

It follows that

$$\varepsilon_v d_v + \varepsilon_w (s-3)d_w = \varepsilon_w (s-1)d_w \iff \varepsilon_v d_v = 2\varepsilon_w d_w.$$

This completes the proof. \square

For $n \geq 3$, it is easy to see from Proposition 5.3 and Proposition 5.4 that if the simplex $\Delta(j, \varepsilon, s)$ is (u, v) -antisymmetric, then there is at least one element among the elements $\varepsilon_i d_i$, for all $1 \leq i \leq n$, and $\varepsilon_j d_j - \varepsilon_i d_i$, for all $1 \leq i < j \leq n$, which is non invertible and equal to zero in $\mathbb{Z}/m\mathbb{Z}$. In this case, the hypothesis of Theorem 3.2 are not satisfied. Therefore, in the next subsection, we only consider the case of dimension $n = 2$.

5.2. In dimension 2. An arithmetic array of dimension 1 is simply called an *arithmetic progression* and is denoted by $\text{AP}(a, d)$ or $\text{AP}(a, d, s)$, for an arithmetic progression of length s , that is,

$$\text{AP}(a, d, s) = (a, a + d, a + 2d, \dots, a + (s - 1)d).$$

Let $W = (w_{-r}, \dots, w_r) \in \mathbb{Z}^{2r+1}$ be the weight sequence of the ACA of dimension 1 that we consider here. By Lemma 3.10, we know that the derived sequence of an arithmetic progression is also an arithmetic progression. Indeed, we have

$$\partial \text{AP}(a, d) = \text{AP}(\sigma a + \sigma' d, \sigma d),$$

where

$$\sigma = \sum_{i=-r}^r w_i \quad \text{and} \quad \sigma' = \sum_{i=-r}^r i w_i.$$

As already remarked, for $\overline{W} = (0, \sigma - \sigma', \sigma')$, we obtain that $\sigma(\overline{W}) = \sigma(W)$ and $\sigma'(\overline{W}) = \sigma'(W)$. Thus, the orbits of $\text{AP}(a, d)$ are the same if we consider W or \overline{W} . Therefore, in the sequel of this subsection, we only consider the case where $r = 1$ and $W = (0, \sigma - \sigma', \sigma')$.

5.2.1. (1,2)-antisymmetric triangles. First, we know from Proposition 5.3 that if the triangle $\Delta(j, \varepsilon, s)$, appearing in the orbit $\mathcal{O}(\text{AP}(a, d))$ with d invertible, is $(1, 2)$ -antisymmetric, then

$$\varepsilon_1 d + \varepsilon_2 \sigma^{-1} \sigma' d = 0 \iff \varepsilon_1 \sigma + \varepsilon_2 \sigma' = 0 \iff \sigma' = -\varepsilon_1 \varepsilon_2 \sigma.$$

So, we deduce that

$$W = (0, (1 + \varepsilon_1 \varepsilon_2) \sigma, -\varepsilon_1 \varepsilon_2 \sigma).$$

Since $\Delta(j, \varepsilon, s)$ is $(1, 2)$ -antisymmetric, we know that $a_j = 0$ and $a_{j+\varepsilon \cdot e_1} + a_{j+\varepsilon \cdot e_2} = 0$. It follows that

$$a_j = 0 \iff a + j_1 d_1 + j_2 d_2 = 0,$$

and

$$\begin{aligned} a_{j+\varepsilon \cdot e_1} + a_{j+\varepsilon \cdot e_2} = 0 &\iff \sigma^{j_2} (a + (j_1 + \varepsilon_1) d_1 + j_2 d_2) + \sigma^{j_2 + \varepsilon_2} (a + j_1 d_1 + (j_2 + \varepsilon_2) d_2) = 0 \\ &\implies \varepsilon_1 d_1 + \sigma^{\varepsilon_2} \varepsilon_2 d_2 = 0 \\ &\iff \varepsilon_1 + \sigma^{\varepsilon_2} \varepsilon_2 \sigma^{-1} \sigma' = 0. \end{aligned}$$

This implies that $\sigma^{\varepsilon_2} = 1$ and thus $\sigma = 1$. Therefore,

$$W = (0, 1 + \varepsilon_1\varepsilon_2, -\varepsilon_1\varepsilon_2).$$

Finally, since $\sigma = 1$, we know that $\Delta(j, \varepsilon, s)$ is an arithmetic triangle which is already balanced for all $s \equiv 0$ or $-1 \pmod m$ from Theorem 3.2.

5.2.2. $(0,2)$ -antisymmetric triangles. First, we know from Proposition 5.4 that if the triangle $\Delta(j, \varepsilon, s)$, appearing in the orbit $\mathcal{O}(\text{AP}(a, d))$ with d invertible, is $(0, 2)$ -antisymmetric, then

$$\varepsilon_2\sigma^{-1}\sigma' = 2\varepsilon_1 \iff \sigma' = 2\varepsilon_1\varepsilon_2\sigma.$$

So, we deduce that

$$W = (0, (1 - 2\varepsilon_1\varepsilon_2)\sigma, 2\varepsilon_1\varepsilon_2\sigma).$$

Since $\Delta(j, \varepsilon, s)$ is $(0, 2)$ -antisymmetric, we know that $a_{j+(s-1)\varepsilon\cdot e_1} = 0$ and $a_{j+(s-2)\varepsilon\cdot e_1} + a_{j+\varepsilon\cdot((s-2)e_1+e_2)} = 0$. It follows that

$$a_{j+(s-1)\varepsilon\cdot e_1} = 0 \iff a + (j_1 + \varepsilon_1(s-1))d_1 + j_2d_2 = 0,$$

and

$$\begin{aligned} a_{j+(s-2)\varepsilon\cdot e_1} + a_{j+\varepsilon\cdot((s-2)e_1+e_2)} = 0 &\iff \sigma^{j_2}(a + (j_1 + \varepsilon_1(s-2))d_1 + j_2d_2) \\ &\quad + \sigma^{j_2+\varepsilon_2}(a + (j_1 + \varepsilon_1(s-2))d_1 + (j_2 + \varepsilon_2)d_2) = 0 \\ &\implies -\varepsilon_1d_1 + \sigma^{\varepsilon_2}(-\varepsilon_1d_1 + \varepsilon_2d_2) = 0 \\ &\iff -\varepsilon_1 + \sigma^{\varepsilon_2}(-\varepsilon_1 + \varepsilon_2\sigma^{-1}\sigma') = 0 \\ &\iff \sigma' = \frac{1 + \sigma^{\varepsilon_2}}{\sigma^{\varepsilon_2}}\varepsilon_1\varepsilon_2\sigma \end{aligned}$$

This implies that $\sigma^{\varepsilon_2} = 1$ and thus $\sigma = 1$. Therefore,

$$W = (0, 1 - 2\varepsilon_1\varepsilon_2, 2\varepsilon_1\varepsilon_2).$$

Finally, since $\sigma = 1$, we know that $\Delta(j, \varepsilon, s)$ is an arithmetic triangle which is already balanced for all $s \equiv 0$ or $-1 \pmod m$ from Theorem 3.2.

5.2.3. $(0,1)$ -antisymmetric triangles. First, we know from Proposition 5.4 that if the simplex $\Delta(j, \varepsilon, s)$ is $(0, 1)$ -antisymmetric, then

$$\varepsilon_1 = 2\varepsilon_2\sigma^{-1}\sigma' \iff \sigma = 2\varepsilon_1\varepsilon_2\sigma'.$$


So, we deduce that

$$W = (0, (2\varepsilon_1\varepsilon_2 - 1)\sigma', \sigma').$$

Now, we refine Theorem 4.1 in this case by considering triangles that have the additional property to be $(0, 1)$ -antisymmetric.

Theorem 5.5. *Let m be an odd positive integers and let $W \in \mathbb{Z}^{2r+1}$ such that $\sigma = 2\sigma'$ and σ is invertible modulo m . Let $a, d \in \mathbb{Z}/m\mathbb{Z}$ such that d is invertible. Then, in the orbit $\mathcal{O}(\text{AP}(a, d))$, every $(0, 1)$ -antisymmetric triangle of orientation $(++)$ or $(--)$ and of size s is balanced, for all positive integers $s \equiv 0$ or $-1 \pmod{\text{lcm}(\text{pord}_m(\sigma), m)}$, where $\text{pord}_m(\sigma)$ is the multiplicative order of σ in $(\mathbb{Z}/m\mathbb{Z})^* / \{-1, 1\}$.*

Proof. Let $\Delta = \Delta(j, \varepsilon, s)$ be a triangle of size s and with orientation $\varepsilon = ++$ or $--$ appearing in the orbit $\mathcal{O}(\text{AP}(a, d)) = (a_i)_{i \in \mathbb{Z} \times \mathbb{N}}$. It is clear that $\text{ord}_m(\sigma) = \text{pord}_m(\sigma)$ or $\text{ord}_m(\sigma) = 2\text{pord}_m(\sigma)$. From Theorem 4.1, we already know that Δ is balanced for $s \equiv 0$ or $-1 \pmod{\text{lcm}(\text{ord}_m(\sigma), m)}$. Suppose now that $\text{ord}_m(\sigma) = 2\text{pord}_m(\sigma)$. Different cases on the congruence of s modulo $\text{pord}_m(\sigma)$ are distinguished.

FIGURE 10. Decomposition of $\Delta' = \Delta(j, \varepsilon, 2s)$

Case 1. $s \equiv 0 \pmod{\text{lcm}(\text{pord}_m(\sigma), m)}$.

We consider the triangle $\Delta' = \Delta(j, \varepsilon, 2s)$. Since $2s \equiv 0 \pmod{\text{lcm}(\text{ord}_m(\sigma), m)}$, the triangle Δ' is balanced by Theorem 4.1. As depicted in Figure, Δ' can be decomposed into two triangles of size s , that are $\Delta(j, \varepsilon, s)$ and $\Delta((j_1, j_2 + \varepsilon_2s), \varepsilon, s)$, and s arithmetic progressions of length s , that are

$$S_k := \{a_{j_1 + \varepsilon_1 l, j_2 + \varepsilon_2 k} \mid l \in [s - k, 2s - k - 1]\},$$

for all $k \in [0, s - 1]$. Since

$$\partial^{j_2 + \varepsilon_2 k} \text{AP}(a, d) = \text{AP}(\sigma^{j_2 + \varepsilon_2 k}(a + (j_2 + \varepsilon_2 k)\sigma^{-1}\sigma'd), \sigma^{j_2 + \varepsilon_2 k}d),$$

for all $k \in [0, s - 1]$, by Proposition 2.7, we deduce that S_k is an arithmetic progression with invertible common difference and of length s , which is divisible by m . Therefore the sequence S_k is balanced in $\mathbb{Z}/m\mathbb{Z}$ for all $k \in [0, s - 1]$. Moreover, since $s \equiv 0 \pmod{\text{lcm}(\text{pord}_m(\sigma), m)}$, we obtain that $\partial^{j_2 + \varepsilon_2 s} \text{AP}(a, d) = \pm \partial^{j_2} \text{AP}(a, d)$. If $\partial^{j_2 + \varepsilon_2 s} \text{AP}(a, d) = \partial^{j_2} \text{AP}(a, d)$, then $\Delta((j_1, j_2 + \varepsilon_2s), \varepsilon, s) = \Delta(j, \varepsilon, s)$ and two copies of Δ can then be seen as the multiset difference of the balanced triangle Δ' and all the arithmetic progressions S_k , which are also balanced. Therefore, Δ is balanced in this case. Otherwise, if $\partial^{j_2 + \varepsilon_2 s} \text{AP}(a, d) = -\partial^{j_2} \text{AP}(a, d)$, then $\Delta((j_1, j_2 + \varepsilon_2s), \varepsilon, s)$ is the opposite triangle of Δ . Moreover, since Δ is antisymmetric, it follows that

$$\mathbf{m}_{\Delta((j_1, j_2 + \varepsilon_2s), \varepsilon, s)}(x) = \mathbf{m}_{\Delta}(-x) = \mathbf{m}_{\Delta}(x)$$

for all $x \in \mathbb{Z}/m\mathbb{Z}$. Finally, since $\Delta((j_1, j_2 + \varepsilon_2s), \varepsilon, s)$ and Δ have the same multiplicity function and since they can be seen as the multiset difference of the balanced triangle Δ' and all the arithmetic progressions S_k , which are balanced, we deduce that Δ is also balanced in this case.

Case 2. $s \equiv -1 \pmod{\text{lcm}(\text{pord}_m(\sigma), m)}$.

The triangle $\Delta(j, \varepsilon, s)$ can be seen as the multiset difference of $\Delta((j_1, j_2 + \varepsilon_2), \varepsilon, s + 1)$, which is balanced by Case 1, and the arithmetic progression $\text{AP}(a_{j_1, j_2 + \varepsilon_2}, \sigma^{j_2 + \varepsilon_2}d, s + 1)$ of invertible common difference and of length $s + 1 \equiv 0 \pmod{m}$, which is also balanced. The multiset difference of two balanced multisets is balanced. This completes the proof. \square

Theorem 5.6. *Let m be an odd positive integers and let $W \in \mathbb{Z}^{2r+1}$ such that $\sigma = -2\sigma'$ and σ is invertible modulo m . Let $a, d \in \mathbb{Z}/m\mathbb{Z}$ such that d is invertible. Then, in the orbit $\mathcal{O}(\text{AP}(a, d))$, every $(0, 1)$ -antisymmetric triangle of orientation $(-+)$ or $(+-)$ and of size s is balanced, for all positive integers $s \equiv 0$ or $-1 \pmod{\text{lcm}(\text{pord}_m(\sigma), m)}$, where $\text{pord}_m(\sigma)$ is the multiplicative order of σ in $(\mathbb{Z}/m\mathbb{Z})^* / \{-1, 1\}$.*

Proof. Similar to the proof of Theorem 5.5 \square

6. CONCLUSION AND OPEN PROBLEMS

Throughout this paper, the existence of infinitely many balanced simplices of $\mathbb{Z}/m\mathbb{Z}$ appearing in the orbit generated from arithmetic arrays by additive cellular automata have been shown and this, for an infinite number of values m for each additive cellular automata of dimension 1 or higher.

For the Pascal cellular automaton of dimension 1, we have proved the existence of infinitely many balanced triangles with orientation $+ -$ and $- +$ in any $\mathbb{Z}/m\mathbb{Z}$ with m odd and with orientation $++$ and $--$ in any $\mathbb{Z}/m\mathbb{Z}$ with m odd not divisible by 3. The following problems remain open.

Problem 6.1. For m even, does there exist infinitely many balanced triangles of $\mathbb{Z}/m\mathbb{Z}$, with any orientation, generated by PCA_1 ?

Problem 6.2. For m odd divisible by 3, does there exist infinitely many balanced triangles of $\mathbb{Z}/m\mathbb{Z}$, with orientations $++$ and $--$, generated by PCA_1 ?

For the Pascal cellular automaton of dimension 2, we have proved the existence of infinitely many balanced tetrahedra with orientation $++-$, $+ - +$, $+ - -$, $- + +$, $- + -$ and $- - -$ in any $\mathbb{Z}/m\mathbb{Z}$ with m odd not divisible by 3 or 5, or m even not divisible by 3 such that $v_2(m) = 1$ and with orientation $++-$ and $- - +$ in any $\mathbb{Z}/m\mathbb{Z}$ with m not divisible by 3 or 5 such that $v_2(m) \leq 1$. The following problems remain open.

Problem 6.3. For m divisible by 3, does there exist infinitely many balanced tetrahedra of $\mathbb{Z}/m\mathbb{Z}$, with any orientation, generated by PCA_2 ?

Problem 6.4. For m even such that $v_2(m) \geq 2$, does there exist infinitely many balanced tetrahedra of $\mathbb{Z}/m\mathbb{Z}$, with any orientation, generated by PCA_2 ?

Problem 6.5. For m odd divisible by 5, does there exist infinitely many balanced tetrahedra of $\mathbb{Z}/m\mathbb{Z}$, with orientations $++-$, $+ - +$, $+ - -$, $- + +$, $- + -$ and $- - -$, generated by PCA_2 ?

Problem 6.6. For m divisible by 5 such that $v_2(m) \leq 1$, does there exist infinitely many balanced tetrahedra of $\mathbb{Z}/m\mathbb{Z}$, with orientations $++-$ and $- - +$, generated by PCA_2 ?

REFERENCES

- [1] M. Augier and S. Eliahou. Parity-regular Steinhaus graphs. *Math. Comp.*, 77(263):1831–1839, 2008.
- [2] C. Bailey and W. Dymacek. Regular Steinhaus graphs. *Congr. Numer.*, 66:45–47, 1988. Nineteenth Southeastern Conference on Combinatorics, Graph Theory, and Computing (Baton Rouge, LA, 1988).
- [3] A. Barbé. Symmetric patterns in the cellular automaton that generates Pascal’s triangle modulo 2. *Discrete Appl. Math.*, 105(1-3):1–38, 2000.
- [4] M. Bartsch. *Steinhaus figures modulo 2 and generalized Steinhaus triangles. (Steinhaus-Figuren modulo 2 und verallgemeinerte Steinhaus-Dreiecke.)*. PhD thesis, Naturwissenschaftliche Fakultät der Technischen Universität Carolo-Wilhelmina zu Braunschweig. 89 S. , 1985.
- [5] J. M. Brunat and M. Maureso. Symmetries in Steinhaus triangles and in generalized Pascal triangles. *Integers*, 11:#A1, 2011.
- [6] G. J. Chang. Binary triangles. *Bull. Inst. Math. Acad. Sinica*, 11(2):209–225, 1983.
- [7] G. J. Chang, B. DasGupta, W. M. Dymáček, M. Fürer, M. Koerlin, Y.-S. Lee, and T. Whaley. Characterizations of bipartite Steinhaus graphs. *Discrete Math.*, 199(1-3):11–25, 1999.
- [8] J. Chappelon. *Graphes de Steinhaus réguliers et triangles de Steinhaus dans les groupes cycliques*. Thèse de doctorat, Université du Littoral Côte d’Opale, 2008.
- [9] J. Chappelon. On a problem of Molluzzo concerning Steinhaus triangles in finite cyclic groups. *Integers*, 8(1):A37, 2008.
- [10] J. Chappelon. Regular Steinhaus graphs of odd degree. *Discrete Math.*, 309(13):4545–4554, 2009.
- [11] J. Chappelon. On the multiplicative order of a^n modulo n . *J. Integer Seq.*, 13(2):Article 10.2.1, 2010.

- [12] J. Chappelon. A universal sequence of integers generating balanced steinhaus figures modulo an odd number. *J. Combin. Theory Ser. A*, 118(1):291–315, 2011.
- [13] J. Chappelon and S. Eliahou. On the problem of molluzzo for the modulus 4. *Integers*, 12(4):723–739, 2012.
- [14] W. M. Dymacek. Steinhaus graphs. In *Proceedings of the Tenth Southeastern Conference on Combinatorics, Graph Theory and Computing (Florida Atlantic Univ., Boca Raton, Fla., 1979)*, Congress. Numer., XXIII–XXIV, pages 399–412, Winnipeg, Man., 1979. Utilitas Math.
- [15] W. M. Dymacek. Bipartite Steinhaus graphs. *Discrete Math.*, 59(1-2):9–20, 1986.
- [16] W. M. Dymacek, M. Koerlin, and T. Whaley. A survey of Steinhaus graphs. In *Combinatorics, graph theory, and algorithms, Vol. I, II (Kalamazoo, MI, 1996)*, pages 313–323. New Issues Press, Kalamazoo, MI, 1999.
- [17] W. M. Dymacek, J.-G. Speton, and T. Whaley. Planar Steinhaus graphs. In *Proceedings of the Thirty-first Southeastern International Conference on Combinatorics, Graph Theory and Computing (Boca Raton, FL, 2000)*, volume 144, pages 193–206, 2000.
- [18] W. M. Dymacek and T. Whaley. Generating strings for bipartite Steinhaus graphs. *Discrete Math.*, 141(1-3):95–107, 1995.
- [19] S. Eliahou and T. Gensane. *in preparation*, 2014.
- [20] S. Eliahou and D. Hachez. On a problem of Steinhaus concerning binary sequences. *Experiment. Math.*, 13(2):215–229, 2004.
- [21] S. Eliahou and D. Hachez. On symmetric and antisymmetric balanced binary sequences. *Integers*, 5(1):#A6, 2005.
- [22] S. Eliahou, J. M. Marín, and M. P. Revuelta. Zero-sum balanced binary sequences. *Integers*, 7(2):#A11, 2007.
- [23] H. Harborth. Solution of Steinhaus’s problem with plus and minus signs. *J. Combinatorial Theory Ser. A*, 12:253–259, 1972.
- [24] H. Harborth and G. Hurlbert. On the number of ones in general binary Pascal triangles. *J. Combin. Math. Combin. Comput.*, 54:99–110, 2005.
- [25] J. C. Molluzzo. Steinhaus graphs. In *Theory and applications of graphs (Proc. Internat. Conf., Western Mich. Univ., Kalamazoo, Mich., 1976)*, volume 642 of *Lecture Notes in Math.*, pages 394–402. Springer, Berlin, 1978.
- [26] H. Steinhaus. *One hundred problems in elementary mathematics*. Basic Books Inc. Publishers, New York, 1964. With a foreword by Martin Gardner.

UNIVERSITÉ MONTPELLIER 2, INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER, CASE COURRIER 051, PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 05, FRANCE
E-mail address: jonathan.chappelon@um2.fr
URL: <http://www.math.univ-montp2.fr/~chappelon/>